

Twisted Ruelle zeta function and complex-valued analytic torsion

Polyxeni Spilioti

April 29, 2020

Abstract

In this paper, we study the twisted Ruelle zeta function associated with the geodesic flow of a compact, hyperbolic, odd-dimensional manifold X . The twisted Ruelle zeta function is associated with an acyclic representation $\chi: \pi_1(X) \rightarrow \mathrm{GL}_n(\mathbb{C})$, which is close enough to an acyclic, unitary representation. In this case, the twisted Ruelle zeta function is regular at zero and equals the square of the refined analytic torsion, as it is introduced in [BK08b], multiplied by an exponential, which involves the eta invariant of the even part of the odd-signature operator associated with χ .

Keywords — Twisted Ruelle zeta function, determinant formula, Cappell-Miller torsion, refined analytic torsion.

Contents

1	Introduction	2
2	Preliminaries	8
2.1	Locally symmetric spaces and representation theory of Lie groups	8
2.1.1	Compact hyperbolic manifolds	8
2.1.2	Representation theory of Lie groups	8
2.1.3	The Casimir element	9
2.2	Preliminaries on the Eta invariant of a non self-adjoint operator and graded regularized determinants	9
2.2.1	Eta invariant of an elliptic operator	9
2.2.2	Regularized determinant of an elliptic operator	10
2.2.3	Graded regularized determinant of an elliptic operator	11
3	Twisted Selberg and Ruelle zeta functions	11
4	Harmonic analysis on symmetric spaces	13
4.1	Twisted Bochner-Laplace operator	13
4.2	The twisted operator $A_\chi^\sharp(\sigma)$	15
5	The determinant formula	16

6	Spaces of representations of the fundamental group	18
6.1	Odd signature operator	18
6.2	Spaces of connections	18
6.3	Spaces of representations	19
7	Refined analytic torsion	20
7.1	Definition	20
7.2	Independence of the refined analytic torsion of the Riemannian metric and the Agmon angle	21
8	Alternative definition of the refined analytic torsion	22
9	Cappell-Miller complex torsion	23
10	Ruelle zeta function and refined analytic torsion	24
11	Appendix: Refined analytic torsion as an element of the determinant line	26
11.1	Refined torsion of a finite-dimensional complex with a chirality operator	27
11.2	Refined analytic torsion as an element of the determinant line for compact odd-dimensional manifolds	29

1 Introduction

The twisted dynamical zeta functions of Ruelle and Selberg are dynamical zeta functions, which are associated with the geodesic flow on the unit sphere bundle $S(X)$ of a compact, hyperbolic manifold X . They are defined in terms of the lengths of the closed geodesics, also called length spectrum. The twisted dynamical zeta functions are defined by Euler-type products, which converge in some right half-plane of \mathbb{C} . The main goal of this paper is to prove that the twisted Ruelle zeta function associated with an acyclic representation of the fundamental group of X , which is close enough to an acyclic and unitary one, is regular at zero and moreover is equal to the Cappell-Miller torsion as it is introduced in [CM10]. Since for such representations, there exists a precise relation between the Cappell-Miller torsion and the refined analytic torsion, as it is introduced in [BK08b], we conclude that twisted Ruelle zeta function at zero is equal to the square of the refined analytic torsion multiplied by an exponential, which involves the eta invariant of the even part of the odd-signature operator associated with χ .

We consider an oriented, compact, hyperbolic manifold X of odd dimension d , obtained as follows. Let either $G = \mathrm{SO}^0(d, 1)$, $K = \mathrm{SO}(d)$ or $G = \mathrm{Spin}(d, 1)$, $K = \mathrm{Spin}(d)$. Then, K is a maximal compact subgroup of G . Let $\tilde{X} := G/K$. The space \tilde{X} can be equipped with a G -invariant metric, which is unique up to scaling and is of constant negative curvature. If we normalize this metric such that it has constant negative curvature -1 , then \tilde{X} , equipped with this metric, is isometric to the d -dimensional hyperbolic space \mathbb{H}^d . Let Γ be a discrete torsion-free subgroup of G such that $\Gamma \backslash G$ is compact. Then, Γ acts by isometries on \tilde{X} and $X = \Gamma \backslash \tilde{X}$ is a compact, oriented, hyperbolic manifold of dimension d .

This is a case of a locally symmetric space of non-compact type of real rank 1. This means that in the Iwasawa decomposition $G = KAN$, A is a multiplicative torus of dimension 1, i.e., $A \cong \mathbb{R}^+$.

For a given $\gamma \in \Gamma$ we denote by $[\gamma]$ the Γ -conjugacy class of γ . If $\gamma \neq e$, then there is a unique closed geodesic c_γ associated with $[\gamma]$. Let $l(\gamma)$ denote the length of c_γ . The conjugacy class $[\gamma]$ is called primitive if there exists no $k \in \mathbb{N}$ with $k > 1$ and $\gamma_0 \in \Gamma$ such that $\gamma = \gamma_0^k$. The prime closed geodesics correspond to the primitive conjugacy classes and are those geodesics that trace out their image exactly once. Let M be the centralizer of A in K . Since Γ is a cocompact subgroup of G , every element $\gamma \in \Gamma - \{e\}$ is hyperbolic. Then, by [Wal76, Lemma 6.5], there exist a $g \in G$, a $m_\gamma \in M$, and an $a_\gamma \in A$, such that $g^{-1}\gamma g = m_\gamma a_\gamma$. The element m_γ is determined up to conjugacy in M , and the element a_γ depends only on γ .

Let $\mathfrak{g}, \mathfrak{a}$ be the Lie algebras of G and A , respectively. Let $\Delta^+(\mathfrak{g}, \mathfrak{a})$ be the set of positive roots of $(\mathfrak{g}, \mathfrak{a})$. Then, $\Delta^+(\mathfrak{g}, \mathfrak{a})$ consists of a single root α . Let \mathfrak{g}_α be the corresponding root space. Let $\bar{\mathfrak{n}}$ be the negative root space of $(\mathfrak{g}, \mathfrak{a})$. Let $S^k(\text{Ad}(m_\gamma a_\gamma)|_{\bar{\mathfrak{n}}})$ be the k -th symmetric power of the adjoint map $\text{Ad}(m_\gamma a_\gamma)$ restricted to $\bar{\mathfrak{n}}$ and ρ be defined as $\rho := \frac{1}{2} \dim(\mathfrak{g}_\alpha)\alpha$. We define the twisted zeta functions associated with unitary, irreducible representations σ of M and finite-dimensional, complex representations χ of Γ . The twisted Selberg zeta function $Z(s; \sigma, \chi)$ is defined for $s \in \mathbb{C}$ by the infinite product

$$Z(s; \sigma, \chi) := \prod_{\substack{[\gamma] \neq e, \\ [\gamma] \text{ prime}}} \prod_{k=0}^{\infty} \det \left(\text{Id} - (\chi(\gamma) \otimes \sigma(m_\gamma) \otimes S^k(\text{Ad}(m_\gamma a_\gamma)|_{\bar{\mathfrak{n}}})) e^{-(s+|\rho|)l(\gamma)} \right). \quad (1.1)$$

The twisted Ruelle zeta function $R(s; \sigma, \chi)$ is defined for $s \in \mathbb{C}$ by the infinite product

$$R(s; \sigma, \chi) := \prod_{\substack{[\gamma] \neq e \\ [\gamma] \text{ prime}}} \det(\text{Id} - \chi(\gamma) \otimes \sigma(m_\gamma) e^{-sl(\gamma)}). \quad (1.2)$$

Both $Z(s; \sigma, \chi)$ and $R(s; \sigma, \chi)$ converge absolutely and uniformly on compact subsets of some half-plane of \mathbb{C} ([Spi18, Proposition 3.4 and Proposition 3.5]).

Fried in [Fri86] proved that for a closed, oriented, hyperbolic manifold $X = \Gamma \backslash \mathbb{H}^d$, and an acyclic, orthogonal representation ϱ_1 of $\pi_1(S(X))$, the Ruelle zeta function, defined for $\text{Re}(s) > d - 1$ by

$$R(s; \varrho_1) := \prod_{\substack{[\gamma] \neq e, \\ [\gamma] \text{ prime}}} \det(\text{Id} - \varrho_1(\gamma) e^{-sl(\gamma)}),$$

admits a meromorphic extension to \mathbb{C} , and for $\epsilon = (-1)^{d-1}$

$$|R(0; \varrho)^\epsilon| = \tau_\varrho(S(X)), \quad (1.3)$$

where $\tau_{\varrho_1}(S(X)) \in \mathbb{R}^+$ is the Reidemeister torsion, which is a topological invariant ([Rei35], [Fra35], [de 36]).

Fried in [Fri87] conjectured that the same result holds for all compact locally homogeneous Riemannian spaces X and acyclic bundles over $S(X)$. This conjecture was proved by [MS91] and [She18] for negatively curved locally symmetric spaces.

Fried, in the same paper, dealt mostly with hyperbolic manifolds of odd dimension. Considering an acyclic, orthogonal representation ϱ of $\pi(X)$ and using the Selberg trace formula for the heat operator $e^{-t\Delta_j}$, where Δ_j is the Hodge Laplacian on j -forms on X with values in the flat vector bundle E_ϱ associated with ϱ , he proved the meromorphic continuation of the zeta functions to the whole complex plane, as well as functional equations for the Selberg zeta function ([Fri86, p. 531-532]). In addition, he proved that $R(0; \varrho)$ is regular at $s = 0$ and

$$R(0; \varrho) = T_\varrho(X)^2, \quad (1.4)$$

where $T_\varrho(X)$ is the Ray-Singer analytic torsion defined as in [RS71] (see [Fri86, eq. (18)]). By Cheeger-Müller theorem ([Che79], [Mül78]), the analytic torsion $T_\varrho(X)$ is equal to the Reidemeister torsion $\tau_\varrho(X)$. Since $\pi_1(S(X)) \cong \pi_1(X)$ and $\tau_\varrho(X)^2 = \tau_{\varrho_1}(S(X))$ ([Fri86, p. 526]), one obtains (1.3). In fact, Fried proved a more general result. If the representation ϱ is not assumed acyclic, then by [Fri86, Theorem 3], the leading term in the Laurent expansion of $R(s; \varrho)$ at $s = 0$ is

$$C_\varrho T_X(\varrho)^2 s^e,$$

where e is a linear combination of the twisted Betti numbers $\beta_i = \dim H^i(X; \varrho)$, and $C_\varrho \in \mathbb{Q}$ is defined in terms of β_i . The theorem of Fried gave rise to important applications, such as the asymptotic behavior of the analytic torsion ([Mül12]) and the study of the growth of the torsion in the cohomology of a closed arithmetic hyperbolic 3-manifold ([MM13]).

The connection of the Ruelle zeta function to spectral invariants such as the analytic torsion has been studied by Bunke and Olbrich ([BO95]), Wotzke ([Wot08]), Müller ([Mül12]) for compact hyperbolic manifolds, under certain assumptions on the representation of the fundamental group of the manifold. For the case of a hyperbolic manifold of finite volume, we refer the reader to the work of Park ([Par09]) and Pfaff ([Pfa14], [Pfa15]). An advanced study of the dynamical zeta functions of locally symmetric manifolds of higher rank is due to Moscovici and Stanton in [MS91], Deitmar in [Dei95], Schen in [She18] and Moscovici, Stanton and Frahm in [MSF18]. We mention also the work of Dang, Guillarmou, Rivier and Shen ([DGRS19]) where the Fried's conjecture is treated for dimension 3, for Anosov flows near the geodesic flow on the unit tangent bundle of hyperbolic 3-manifolds. This result was extended by Chaubet and Dang to higher dimensions in [CD19], where a new object is introduced, the dynamical torsion.

For unitary representations, the dynamical zeta functions have been studied by Bunke and Olbrich in [BO95] for all locally symmetric spaces of real rank 1. They proved that the zeta functions admit a meromorphic continuation to the whole complex plane and satisfy functional equations. Moreover, for compact, hyperbolic, odd dimensional manifolds, and under the additional assumption that the representation is acyclic, they proved that the Ruelle zeta function is regular at zero and equals the square of the analytic torsion as in (1.4) ([BO95, Theorem 4.8]).

Wotzke in [Wot08] extended this result for representations of Γ , which are not necessary unitary, but very special ones. In particular, he considered a compact odd-dimensional, hyperbolic, manifold and a finite-dimensional, irreducible, representation $\tau: G \rightarrow \mathrm{GL}(V)$ of G , such that $\tau \neq \tau_\theta$, where $\tau_\theta = \tau \circ \theta$ and θ denotes the Cartan involution of G . Under these assumptions, he proved

that the Ruelle zeta function admits a meromorphic continuation to the whole complex plane. In addition, it is regular at zero and

$$|R(0; \tau)| = T_\tau(X)^2,$$

where the representation τ restricted to Γ . Wotzke's method is based on the fact that if one considers the restrictions $\tau|_K$ and $\tau|_\Gamma$ of τ to K and Γ , respectively, there is an isomorphism between the locally homogeneous vector bundle E_τ over X , associated with $\tau|_K$, and the flat vector bundle E_{flat} over X associated with $\tau|_\Gamma$. By [MM63, Lemma 3.1], a Hermitian fiber metric in E_τ descends to a fiber metric in E_{flat} . Therefore, one deals with self-adjoint Laplacians and all tools from harmonic analysis on locally symmetric spaces are available.

In [Spi18], it was proved that the twisted dynamical zeta functions associated with a general, finite-dimensional, complex representation of the fundamental group, as in (1.1) and (1.2), admit a meromorphic continuation to the whole complex plane. In [Spi15], the functional equations for them are derived. Moreover, it is shown a determinant formula, which relates the twisted Ruelle zeta function with a finite product of graded regularized determinants of certain, twisted (non-self-adjoint), elliptic differential operators (see [Spi15, Proposition 7.9, case (a)]). The above results extended the results of Bunke and Olbrich to the case of non-unitary twists. This determinant formula is the keypoint to prove our main results, Theorem 1, Theorem 2 and Corollary 1, below.

In the present paper, we consider acyclic representations of Γ , which are non-unitary but closed enough to unitary ones. We provide here some details about these representations. For a compact, oriented, odd-dimensional, Riemannian manifold (X, g) and a complex vector bundle $E \rightarrow X$, equipped with a flat connection ∇ , Braverman and Kappeler in [BK08b] considered the odd signature operator $B = (\nabla, g)$, acting on the space of smooth differential forms $\Lambda^k(X, E)$ on X , with values in E (see Section 6.2 in [BK08b] and Section 6.1 in the present paper). This is a first order, elliptic differential operator, which is in general non-self-adjoint. Suppose that there exists a Hermitian metric on E , which is preserved by ∇ . In such a case, we say that ∇ is a Hermitian connection. Then the operator is (formally) self-adjoint. Hence, if we assume further that ∇ is acyclic, i.e., $\text{Im}(\nabla|_{\Lambda^{k-1}(X, E)}) = \text{Ker}(\nabla|_{\Lambda^k(X, E)})$, for every k , the odd signature operator B is bijective. If there is no Hermitian metric on E , which is preserved by ∇ , then the odd signature operator is no longer self-adjoint. One can assume that ∇ is acyclic, but this does not imply that the odd signature operator has a trivial kernel, since Hodge theory is no longer applicable.

However, by a continuity argument, [BK08b, Proposition 6.8], the following assumptions

Assumption 1. The connection ∇ is acyclic, i.e., for every k , $\text{Im}(\nabla|_{\Lambda^{k-1}(X, E)}) = \text{Ker}(\nabla|_{\Lambda^k(X, E)})$;

Assumption 2. The odd signature operator $B: \Lambda^k(X, E) \rightarrow \Lambda^k(X, E)$ is bijective;

are satisfied for all flat connections in an open neighbourhood, in a suitable C^0 -topology (see Subsection 6.2), of the set of acyclic, Hermitian connections.

Let $\text{Rep}(\pi_1(X), \mathbb{C}^n)$ be the set of all n -dimensional, complex representations of $\pi_1(X)$. This set has a natural structure of a complex algebraic variety (see Subsection 6.3). Each representation $\chi \in \text{Rep}(\pi_1(X), \mathbb{C}^n)$ gives rise to a vector bundle E_χ with a flat connection ∇_χ , whose monodromy is isomorphic to χ .

Let $\text{Rep}_0(\pi_1(X), \mathbb{C}^n)$ be the set of all acyclic representations of $\pi_1(X)$, i.e., the set of representations χ such that ∇_χ is acyclic. Let $\text{Rep}^u(\pi_1(X), \mathbb{C}^n)$ be the set of all unitary representations of $\pi_1(X)$, i.e., the set of representations such that there exists a Hermitian scalar product (\cdot, \cdot) on \mathbb{C}^n , which is preserved by the matrices $\chi(\gamma)$, for all $\gamma \in \pi_1(X)$. We set

$$\text{Rep}_0^u(\pi_1(X), \mathbb{C}^n) := \text{Rep}_0(\pi_1(X), \mathbb{C}^n) \cap \text{Rep}^u(\pi_1(X), \mathbb{C}^n).$$

Let $B_\chi := B(\nabla_\chi, g): \Lambda^k(X, E) \rightarrow \Lambda^k(X, E)$. Let B_χ^{even} be the restriction of B_χ to $\Lambda^{\text{even}}(X, E)$. Suppose that for some representation $\chi_0 \in \text{Rep}_0(\pi_1(X), \mathbb{C}^n)$ the operator B_{χ_0} is bijective. Then, there exists an open neighbourhood (in classical topology) $V \subset \text{Rep}(\pi_1(X), \mathbb{C}^n)$ of the set $\text{Rep}_0^u(\pi_1(X), \mathbb{C}^n)$ of acyclic, unitary representations such that, for all $\chi \in V$, the pair (∇_χ, g) satisfies Assumptions 1 and 2. For such representations Braverman and Kappeler defined a Riemannian metric-invariant, the refined analytic torsion, which is a non zero complex number, defined by

$$T_\chi := \det_{\text{gr}, \theta}(B_\chi^{\text{even}}) e^{i\pi \text{rank}(E_\chi) \eta_{\text{tr}}(g)}.$$

Here $\theta \in (-\pi, 0)$ is an Agmon angle for B_χ^{even} , $\det_{\text{gr}, \theta}(B_\chi^{\text{even}})$ denotes the graded determinant of B_χ^{even} (see Remark 7.1.2), and $\eta_{\text{tr}}(g) = \frac{1}{2} \eta(0, B_{\text{tr}}(g))$, where $\eta(0, B_{\text{tr}}(g))$ denotes the eta invariant of the even part of the odd signature operator corresponding to the trivial line bundle, endowed with the trivial connection (see Definition 7.1.1). One can view this definition as a special case of the definition of the refined analytic torsion as an element of the determinant line in [BK07], where the assumption of the bijectivity of the odd signature operator is removed.

On the other hand, Cappell and Miller in [CM10], defined another invariant, the Cappell-Miller torsion $\tau \in \det(H^*(X, E)) \otimes \det(H^*(X, E))$. In this setting, the regularized determinants of the flat Hodge Laplacians are equipped. To define the flat Laplacians, no use of a Hermitian metric on the flat vector bundle is needed (see Section 9). These operators coincide with the square B^2 of the odd signature operator B . If $\chi \in V$, the element of $\det(H^*(X, E_\chi)) \otimes \det(H^*(X, E_\chi))$ does not contribute in the definition (see Definition 9.2). In such a case, the Cappell-Miller torsion τ_χ is a complex number given by

$$\tau_\chi := \prod_{k=0}^d \det(B_{\chi, k}^2)^{k(-1)^{k+1}}.$$

Then, by Proposition 9.4 (see also [BK08a, Subsection 5.3]), for $\chi \in V$,

$$\tau_\chi = T_\chi^2 e^{2\pi i(\eta(B_\chi^{\text{even}}) - \text{rank}(E_\chi) \eta_{\text{tr}})},$$

where $\eta(B_\chi^{\text{even}})$ denotes the eta invariant of the even part of the odd signature operator B_χ .

Let now X be an oriented, compact, hyperbolic, odd-dimensional manifold $X = \Gamma \backslash \mathbb{H}^d$ as above. By the determinant formula in [Spi15], we obtain the following theorem.

Theorem 1. *Let χ be a finite-dimensional complex representation of Γ . Let $\Delta_{\chi, k}^\#$ be the flat Hodge Laplacian, acting on the space of k -differential forms on*

X with values in the flat vector bundle E_χ . Then, the Ruelle zeta function has the representation

$$R(s; \chi) = \prod_{k=0}^{d-1} \prod_{p=k}^{d-1} \det_{\text{gr}} (\Delta_{\chi,k}^\# + s(s + 2(|\rho| - p)))^{(-1)^p} \\ \cdot \exp \left((-1)^{\frac{d-1}{2}+1} \pi(d+1) \dim(V_\chi) \frac{\text{Vol}(X)}{\text{Vol}(S^d)} s \right),$$

where $\text{Vol}(S^d)$ denotes the volume of the d -dimensional Euclidean unit sphere. Let $d_{\chi,k} := \dim \text{Ker}(\Delta_{\chi,k})$. Then, the singularity of the Ruelle zeta function at $s = 0$ is of order

$$\sum_{k=0}^{(d-1)/2} (d+1-2k)(-1)^k d_{\chi,k}.$$

Let $V \subset \text{Rep}(\pi_1(X), \mathbb{C}^n)$ be an open neighbourhood (in classical topology) of the set $\text{Rep}_0^u(\pi_1(X), \mathbb{C}^n)$ of acyclic, unitary representations such that, for all $\chi \in V$, B_χ is bijective. Then, for $\chi \in V$, we obtain the following result.

Theorem 2. *Let $\chi \in V$. Then, the Ruelle zeta function $R(s; \chi)$ is regular at $s = 0$ and is equal to the complex Cappell-Miller torsion,*

$$R(0; \chi) = \tau_\chi.$$

Corollary 1. *Let $\chi \in V$. Then the Ruelle zeta function $R(s; \chi)$ is regular at $s = 0$ and is related to the refined analytic torsion T_χ by*

$$R(0; \chi) = T_\chi^2 e^{2\pi i(\eta(B_\chi^{\text{even}}) - \text{rank}(E_\chi)\eta_{\text{tr}})}.$$

Organization of the paper This paper is organised as follows. In Section 2, we summarize basic facts about hyperbolic manifolds, and representations of the involved Lie groups. In addition, we provide the definition of the eta invariant and the regularized determinant of a non-self-adjoint elliptic differential operator. In Section 3, we consider the twisted Selberg and Ruelle zeta functions. In Section 4, we recall the definition of the twisted Laplace operator from [Mül11] and certain auxiliary operators, first introduced in [BO95]. In Section 5, we use the determinant formula from [Spi15] to express the twisted Ruelle zeta function in terms of products of graded regularized determinants of twisted Hodge Laplacians on vector bundle-valued differential forms. This formula is the keypoint to study the singularity of the twisted Ruelle zeta function at $s = 0$. In Section 6, we introduce the odd signature operator and consider representations of the fundamental group, which are non-unitary, but closed enough, in classical topology, to acyclic and unitary representations. In Sections 7 and 8, we consider the refined analytic torsion, as it is introduced in [BK08b]. In Section 9, we consider the Cappell-Miller torsion, as it is introduced in [CM10], and its relation to the refined analytic torsion. Finally, in Section 10, we prove the main results, Theorem 2 and Corollary 1. We include also an Appendix, in Section 10, where we recall the general definition of the refined analytic torsion as an element of the determinant line from [BK07].

Acknowledgements The author would like to thank Werner Müller for his helpful suggestions and comments, as well as Maxim Braverman for the insightful discussions. The author wishes to acknowledge also the hospitality of the Department of Mathematics of the University of Tübingen and of the Department of Mathematics of Aarhus University.

2 Preliminaries

2.1 Locally symmetric spaces and representation theory of Lie groups

2.1.1 Compact hyperbolic manifolds

In this paper, we are dealing with odd-dimensional, compact, hyperbolic manifolds, obtained as follows. Let $d = 2n + 1$, $n \in \mathbb{N}_{>0}$. We consider the universal coverings $G = \text{Spin}(d, 1)$ of $\text{SO}^0(d, 1)$ and $K = \text{Spin}(d)$ of $\text{SO}(d)$, respectively. We set $\tilde{X} := G/K$. Let $\mathfrak{g}, \mathfrak{k}$ be the Lie algebras of G and K , respectively. Let Θ be the Cartan involution of G and let θ be the differential of Θ at e , which is the identity element of G . Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition of \mathfrak{g} with respect to θ . There exists a canonical isomorphism $T_{eK} \cong \mathfrak{p}$. Let $B(X, Y)$ be the Killing form on $\mathfrak{g} \times \mathfrak{g}$ defined by $B(X, Y) = \text{Tr}(\text{ad}(X) \circ \text{ad}(Y))$. It is a symmetric bilinear form. We set

$$\langle Y_1, Y_2 \rangle := \frac{1}{2(d-1)} B(Y_1, Y_2), \quad Y_1, Y_2 \in \mathfrak{g}. \quad (2.1)$$

The restriction of $\langle \cdot, \cdot \rangle$ to \mathfrak{p} defines an inner product on \mathfrak{p} and therefore induces a G -invariant Riemannian metric on \tilde{X} , which has constant curvature -1 . Then, \tilde{X} equipped with this metric is isometric to \mathbb{H}^d . Let Γ_1 be a torsion-free, co-compact, discrete subgroup of $\text{SO}^0(d, 1)$. We assume that Γ_1 can be lifted to a subgroup Γ of G . Then, $X := \Gamma \backslash \tilde{X}$ is a compact, hyperbolic manifold of odd dimension d .

2.1.2 Representation theory of Lie groups

Let

$$G = KAN$$

be the standard Iwasawa decomposition of G and let M be the centralizer of A in K . Then, $M = \text{Spin}(d-1)$. Let \mathfrak{a} and \mathfrak{m} be the Lie algebras of A and M , respectively. Let \mathfrak{b} be a Cartan subalgebra of \mathfrak{m} . Let \mathfrak{h} be a Cartan subalgebra of \mathfrak{g} . We consider the complexifications $\mathfrak{g}_{\mathbb{C}} := \mathfrak{g} \oplus i\mathfrak{g}$, $\mathfrak{h}_{\mathbb{C}} := \mathfrak{h} \oplus i\mathfrak{h}$ and $\mathfrak{m}_{\mathbb{C}} := \mathfrak{m} \oplus i\mathfrak{m}$.

Let $\Delta^+(\mathfrak{g}, \mathfrak{a})$ be the set of positive roots of $(\mathfrak{g}, \mathfrak{a})$ and \mathfrak{g}_{α} the corresponding root spaces. In the present case $\Delta^+(\mathfrak{g}, \mathfrak{a})$ consists of a single root α . Let $M' = \text{Norm}_K(A)$ be the normalizer of A in K . Let $H \in \mathfrak{a}$ such that $\alpha(H) = 1$. With respect to the inner product, induced by (2.1) on \mathfrak{a} , H has norm 1. We define

$$A^+ := \{\exp(tH) : t \in \mathbb{R}^+\}; \quad (2.2)$$

$$\rho := \frac{1}{2} \dim(\mathfrak{g}_{\alpha})\alpha; \quad (2.3)$$

$$\rho_{\mathfrak{m}} := \frac{1}{2} \sum_{\alpha \in \Delta^+(\mathfrak{m}_{\mathbb{C}}, \mathfrak{b})} \alpha. \quad (2.4)$$

The inclusion $i: M \hookrightarrow K$ induces the restriction map $i^*: R(K) \rightarrow R(M)$, where $R(K), R(M)$ are the representation rings over \mathbb{Z} of K and M , respectively. Let \widehat{K}, \widehat{M} be the sets of equivalent classes of irreducible unitary representations of K and M , respectively. For the highest weight ν_{τ} of $\tau \in \widehat{K}$, we have $\nu_{\tau} = (\nu_1, \dots, \nu_n)$, where $\nu_1 \geq \dots \geq \nu_n$ and $\nu_i, i = 1, \dots, n$ are all half integers (i.e., $\nu_i = q_i + \frac{1}{2}, q_i \in \mathbb{Z}$). For the highest weight ν_{σ} of $\sigma \in \widehat{M}$, we have

$$\nu_{\sigma} = (\nu_1, \dots, \nu_{n-1}, \nu_n), \quad (2.5)$$

where $\nu_1 \geq \dots \geq \nu_{n-1} \geq |\nu_n|$ and $\nu_i, i = 1, \dots, n$ are all half integers ([BO95, p. 20]). We denote by (s, S) be the spin representation of K and by $(s^+, S^+), (s^-, S^-)$ the half spin representations of M ([Friedrich00, p. 22]).

2.1.3 The Casimir element

Let Z_i be a basis of \mathfrak{g} and let Z^j be the basis of \mathfrak{g} , which is determined by $\langle Z_i, Z^j \rangle = \delta_{ij}$, where $\langle \cdot, \cdot \rangle$ is as in (2.1). Let $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ be the universal enveloping algebra of $\mathfrak{g}_{\mathbb{C}}$ and let $Z(\mathfrak{g}_{\mathbb{C}})$ be its center. Then, $\Omega \in Z(\mathfrak{g}_{\mathbb{C}})$ is given by

$$\Omega = \sum_i Z_i Z^i.$$

By Kuga's Lemma ([MM63, (6.9)]), the Hodge Laplacian, acting on the space $\Lambda^*(G/K)$ of differential forms on G/K , coincides with $-R(\Omega)$, where $R(\Omega)$ is the Casimir operator on $\Lambda^*(G/K)$, induced by Ω .

2.2 Preliminaries on the Eta invariant of a non self-adjoint operator and graded regularized determinants

2.2.1 Eta invariant of an elliptic operator

Let $E \rightarrow X$ be a complex vector bundle over a smooth, compact, Riemannian manifold X of dimension d . Let $D: C^\infty(X, E) \rightarrow C^\infty(X, E)$ be an elliptic differential operator of order $m \geq 1$. Let $\sigma(D)$ be its principal symbol.

Definition 2.2.1. *Let $R_\theta := \{\rho e^{i\theta} : \rho \in [0, \infty)\}$. The angle $\theta \in [0, 2\pi)$ is a principal angle for D if*

$$\text{spec}(\sigma_D(x, \xi)) \cap R_\theta = \emptyset, \quad \forall x \in X, \forall \xi \in T_x^* X, \xi \neq 0.$$

Definition 2.2.2. *Let $I \subset [0, 2\pi)$. Let L_I be a solid angle defined by*

$$L_I := \{\rho e^{i\theta} : \rho \in (0, \infty), \theta \in I\}.$$

The angle θ is an Agmon angle for D , if it is a principal angle for D and there exists an $\varepsilon > 0$ such that

$$\text{spec}(D) \cap L_{[\theta-\varepsilon, \theta+\varepsilon]} = \emptyset.$$

We define here the eta function associated with non-self-adjoint operators D with elliptic, self-adjoint principal symbol (see [Gil84]). By [Mar88, §I.6], the space $L^2(X, E)$ of square integrable sections of E is the closure of the algebraic direct sum of finite dimensional D -invariant subspaces

$$L^2(X, E) = \overline{\bigoplus_{k \geq 1} \Lambda_k},$$

such that the restriction of D to Λ_k has a unique eigenvalue λ_k and $\lim_{k \rightarrow \infty} |\lambda_k| = \infty$. In general, the above sum is not a sum of mutually orthogonal subspaces. The space Λ_k is called the space of root vectors of D with eigenvalue λ_k . The algebraic multiplicity m_k of the the eigenvalue λ_k is defined as the the dimension of the space Λ_k . Since the principal symbol of D is self-adjoint, the angles $\pm\pi/2$ are principal angles for D . Hence, D also possesses an Agmon angle (see [BK08b, Section 3.10]). Let θ be an Agmon angle for D . Denote by $\log_\theta \lambda_k$ the branch of the logarithm in $\mathbb{C} \setminus R_\theta$ with $\theta < \text{Im}(\log_\theta \lambda_k) < \theta + 2\pi$. Let $(\lambda_k)_\theta := e^{\log_\theta \lambda_k}$.

Definition 2.2.3. For $\text{Re}(s) \gg 0$, we define the eta function $\eta_\theta(s, D)$ of D by

$$\eta_\theta(s, D) := \sum_{\text{Re}(\lambda_k) > 0} m_k (\lambda_k)_\theta^{-s} - \sum_{\text{Re}(\lambda_k) < 0} m_k (-\lambda_k)_\theta^{-s}.$$

Note that since the angles $\pm\pi/2$ are principal angles for D , there are at most finitely many eigenvalues of D on or near the imaginary axis. Hence, the eigenvalues of D that are purely imaginary do not contribute to the definition of the eta function. It has been shown by Grubb and Seeley ([GS95, Theorem 2.7]) that $\eta_\theta(s, D)$ has a meromorphic continuation to the whole complex plane \mathbb{C} with isolated simple poles and that is regular at $s = 0$. Moreover, the number $\eta_\theta(0, D)$ is independent of the Agmon angle θ . Hence, we write $\eta(0, D)$ instead of $\eta_\theta(0, D)$.

Definition 2.2.4. Let m_+ , respectively m_- , denote the the number of the eigenvalues of D on the positive, respectively negative, part of the imaginary axis. We define the eta invariant $\eta(D)$ of the operator D by

$$\eta(D) = \frac{\eta(0, D) + m_+ - m_-}{2}.$$

By [BK08b, (3.25)], $\eta(D)$ is independent of the Agmon angle θ .

2.2.2 Regularized determinant of an elliptic operator

We recall here the definition of the regularized determinant of an elliptic operator. For more details we refer the reader to [BK08b, Subsection 3.5 and Definition 3.6]. Let X, E and D be as in the previous subsection. We assume, in addition, that D has a self-adjoint principal symbol and it is invertible. Let θ be an Agmon angle for D . We define the zeta function $\zeta_\theta(s, D)$ by

$$\zeta_\theta(s, D) := \text{Tr}(D_\theta^{-s}), \quad \text{Re}(s) > \frac{\dim(X)}{m},$$

where D_θ^{-s} is a pseudo-differential operator with continuous kernel (see [BK08b, (3.15)]). The zeta function $\zeta_\theta(s, D)$ admits a meromorphic continuation to \mathbb{C}

and it is regular at zero ([See67]). We define the regularized determinant of D by

$$\det_\theta := \exp \left(- \frac{d}{ds} \Big|_{s=0} \zeta_\theta(s, D) \right).$$

We denote by $L\det_\theta$ the particular value of the logarithm of the determinant, such that

$$L\det_\theta(D) = -\zeta'_\theta(0, D).$$

By [BK08b, Subsection 3.10], the regularized determinant $\det_\theta(D)$ is independent of the Agmon angle θ .

2.2.3 Graded regularized determinant of an elliptic operator

We recall here the notion of the graded regularized determinant of an elliptic differential operator. Let $E = E_+ \oplus E_-$ be a \mathbb{Z}_2 -graded vector bundle over a compact, Riemannian manifold X . Let $D: C^\infty(X, E) \rightarrow C^\infty(X, E)$ be an elliptic differential operator, which is bounded from below. We assume that D preserves the grading, i.e., we assume that with respect to the decomposition

$$C^\infty(X, E) = C^\infty(X, E^+) \oplus C^\infty(X, E^-),$$

D takes the form,

$$D = \begin{pmatrix} D_+ & 0 \\ 0 & D_- \end{pmatrix}.$$

Then, the graded determinant $\det_{\text{gr}}(D)$ of D is defined by

$$\det_{\text{gr}}(D) = \frac{\det(D_+)}{\det(D_-)},$$

where $\det(D_+)$ and $\det(D_-)$ denote the regularized determinants of the operators D_+ and D_- , respectively.

3 Twisted Selberg and Ruelle zeta functions

We consider the twisted Ruelle and Selberg zeta functions associated with the geodesic flow on the sphere vector bundle $S(X)$ of $X = \Gamma \backslash G/K$. Since K acts transitively on the unit vectors in \mathfrak{p} , $S(\tilde{X})$ can be represented by the homogeneous space G/M . Therefore $S(X) = \Gamma \backslash G/M$.

We recall the Cartan decomposition $G = KA^+K$ of G , where A^+ is as in (2.2). Then, every element $g \in G$ can be written as $g = ha_+k$, where $h, k \in K$ and $a_+ = \exp(tH)$, for some $t \in \mathbb{R}^+$. The positive real number t equals $d(eK, gK)$, where $d(\cdot, \cdot)$ denotes the geodesic distance on \tilde{X} . It is a well known fact that there is a 1-1 correspondence between the closed geodesics on a manifold X with negative sectional curvature and the non-trivial conjugacy classes of the fundamental group $\pi_1(X)$ of X ([GKM68]).

The hyperbolic elements of Γ can be realized as the semisimple elements of this group, i.e., the diagonalizable elements of Γ . Since Γ is a cocompact subgroup of G , every element $\gamma \in \Gamma - \{e\}$ hyperbolic. We denote by c_γ the closed geodesic on X , associated with the hyperbolic conjugacy class $[\gamma]$. We denote by $l(\gamma)$ the length of c_γ . Since Γ is torsion-free, $l(\gamma)$ is always positive and therefore

we can obtain an infimum for the length spectrum $\text{spec}(\Gamma) := \{l(\gamma) : \gamma \in \Gamma\}$. An element $\gamma \in \Gamma$ is called primitive if there exists no $n \in \mathbb{N}$ with $n > 1$ and $\gamma_0 \in \Gamma$ such that $\gamma = \gamma_0^n$. A primitive element $\gamma_0 \in \Gamma$ corresponds to a geodesic on X . The prime geodesics correspond to the periodic orbits of minimal length. Hence, if a hyperbolic element γ in Γ is generated by a primitive element γ_0 , then there exists a $n_\Gamma(\gamma) \in \mathbb{N}$ such that $\gamma = \gamma_0^{n_\Gamma(\gamma)}$ and the corresponding closed geodesic is of length $l(\gamma) = n_\Gamma(\gamma)l(\gamma_0)$.

We lift now the closed geodesic c_γ to the universal covering \tilde{X} . For $\gamma \in \Gamma$, $l(\gamma) := \inf\{d(x, \gamma x) : x \in \tilde{X}\}$, or $l(\gamma) = \inf\{d(eK, g^{-1}\gamma gK) : g \in G\}$. Hence, we see that the length of the closed geodesic $l(\gamma)$ depends only on $\gamma \in \Gamma$. Let $\gamma \in \Gamma$, with $\gamma \neq e$. Then, by [Wal76, Lemma 6.5] there exist a $g \in G$, a $m_\gamma \in M$, and an $a_\gamma \in A^+$, such that $g^{-1}\gamma g = m_\gamma a_\gamma$. The element m_γ is determined up to conjugacy in M , and the element a_γ depends only on γ . As in [BO95, Section 3.1], we consider the geodesic flow ϕ on $S(X)$, given by the map $\phi: \mathbb{R} \times S(X) \ni (t, \Gamma gM) \rightarrow \Gamma g \exp(-tH)M \in S(X)$. A closed orbit of ϕ is described by the set $c := \{\Gamma g \exp(-tH)M : t \in \mathbb{R}\}$, where $g \in G$ is such that $g^{-1}\gamma g := m_\gamma a_\gamma \in MA^+$. The Anosov property of the geodesic flow ϕ on $S(X)$ can be expressed by the following $d\phi$ -invariant splitting of $TS(X)$

$$TS(X) = T^s S(X) \oplus T^c S(X) \oplus T^u S(X), \quad (3.1)$$

where $T^s S(X)$ consists of vectors that shrink exponentially, $T^u S(X)$ consists of vectors that expand exponentially, and $T^c S(X)$ is the one dimensional subspace of vectors tangent to the flow, with respect to the Riemannian metric, as $t \rightarrow \infty$. The splitting in (3.1) corresponds to the splitting

$$TS(X) = \Gamma \backslash G \times_{\text{Ad}} (\bar{\mathfrak{n}} \oplus \mathfrak{a} \oplus \mathfrak{n}),$$

where Ad denotes the adjoint action of $\text{Ad}(\exp(-tH))$ on $\bar{\mathfrak{n}}, \mathfrak{a}, \mathfrak{n}$, and $\bar{\mathfrak{n}} = \theta \mathfrak{n}$ is the sum of the negative root spaces of the system $(\mathfrak{g}, \mathfrak{a})$.

Definition 3.1. Let $\chi: \Gamma \rightarrow \text{GL}(V_\chi)$ be a finite-dimensional, complex representation of Γ and $\sigma \in \widehat{M}$. The twisted Selberg zeta function $Z(s; \sigma, \chi)$ is defined by the infinite product

$$Z(s; \sigma, \chi) := \prod_{\substack{[\gamma] \neq e \\ [\gamma] \text{ prime}}} \prod_{k=0}^{\infty} \det(\text{Id} - (\chi(\gamma) \otimes \sigma(m_\gamma) \otimes S^k(\text{Ad}(m_\gamma a_\gamma)|_{\bar{\mathfrak{n}}})) e^{-(s+|\rho)|l(\gamma)}),$$

where $s \in \mathbb{C}$, $S^k(\text{Ad}(m_\gamma a_\gamma)|_{\bar{\mathfrak{n}}})$ denotes the k -th symmetric power of the adjoint map $\text{Ad}(m_\gamma a_\gamma)$ restricted to $\bar{\mathfrak{n}}$ and ρ is as in (2.3).

By [Spi18, Proposition 3.4], there exists a positive constant c such that the twisted Selberg zeta function $Z(s; \sigma, \chi)$ converges absolutely and uniformly on compact subsets of the half-plane $\text{Re}(s) > c$.

Definition 3.2. Let $\chi: \Gamma \rightarrow \text{GL}(V_\chi)$ be a finite-dimensional, complex representation of Γ and $\sigma \in \widehat{M}$. The twisted Ruelle zeta function $R(s; \sigma, \chi)$ is defined by the infinite product

$$R(s; \sigma, \chi) := \prod_{\substack{[\gamma] \neq e \\ [\gamma] \text{ prime}}} \det(\text{Id} - (\chi(\gamma) \otimes \sigma(m_\gamma)) e^{-sl(\gamma)})^{(-1)^{d-1}},$$

where $s \in \mathbb{C}$.

By [Spi18, Proposition 3.5], there exists a positive constant r such that the twisted Selberg zeta function $R(s; \sigma, \chi)$ converges absolutely and uniformly on compact subsets of the half-plane $\operatorname{Re}(s) > r$.

The twisted dynamical zeta functions associated with an arbitrary representation of Γ has been studied in [Spi18] and [Spi15]. Specifically, by [Spi18, Theorem 1.1 and Theorem 1.2], the twisted dynamical zeta functions admit a meromorphic continuation to the whole complex plane.

4 Harmonic analysis on symmetric spaces

4.1 Twisted Bochner-Laplace operator

In this section, we define the twisted Bochner-Laplace operator as it is introduced in [Mül11, Section 4]. In addition, we describe this operator in the locally symmetric space setting.

Let $E_0 \rightarrow X$ be a complex vector bundle with covariant derivative ∇ . We define the second covariant derivative ∇^2 by

$$\nabla_{V,W}^2 := \nabla_V \nabla_W - \nabla_{\nabla_V^L C W},$$

where $V, W \in C^\infty(X, TX)$ and ∇^{LC} denotes the Levi-Civita connection on TX . We define the connection Laplacian Δ_{E_0} to be the negative of the trace of the second covariant derivative, i.e.,

$$\Delta_{E_0} := -\operatorname{Tr} \nabla^2.$$

If E_0 is equipped with a Hermitian metric compatible with the connection ∇ , then by [LM89, p. 154], the connection Laplacian is equal to the Bochner-Laplace operator, i.e.,

$$\Delta_{E_0} = \nabla^* \nabla.$$

In terms of a local orthonormal frame field (e_1, \dots, e_d) of $T_x X$, for $x \in X$, the connection Laplacian is given by

$$\Delta_{E_0} = -\sum_{j=1}^d \nabla_{e_j, e_j}^2.$$

The principal symbol of Δ_{E_0} equals

$$\sigma_{\Delta_{E_0}}(x, \xi) = \|\xi\|^2 \operatorname{Id}_{(E_0)_x},$$

where $x \in X$ and $\xi \in T_x^* X$. Δ_{E_0} acts in $L^2(X, E_0)$ with domain $C^\infty(X, E_0)$. The operator $\Delta_{E_0} : C^\infty(X, E_0) \rightarrow C^\infty(X, E_0)$ is a second order, elliptic, formally self-adjoint differential operator.

Let $\chi : \Gamma \rightarrow \operatorname{GL}(V_\chi)$ be a finite-dimensional, complex representation of Γ . Let $E_\chi \rightarrow X$ be the associated flat vector bundle over X , equipped with a flat connection ∇^{E_χ} . We specialize to the twisted case $E = E_0 \otimes E_\chi$, where $E_0 \rightarrow X$ is a complex vector bundle equipped with a connection ∇^{E_0} and a metric, which is compatible with this connection. Let $\nabla^E = \nabla^{E_0 \otimes E_\chi}$ be the product connection, defined by

$$\nabla^{E_0 \otimes E_\chi} := \nabla^{E_0} \otimes 1 + 1 \otimes \nabla^{E_\chi}.$$

We define the operator $\Delta_{E_0, \chi}^\sharp$ by

$$\Delta_{E_0, \chi}^\sharp := -\text{Tr}((\nabla^{E_0 \otimes E_\chi})^2). \quad (4.1)$$

We choose a Hermitian metric in E_χ . Then, $\Delta_{E_0, \chi}^\sharp$ acts on $L^2(X, E_0 \otimes E_\chi)$. However, it is not a formally self-adjoint operator in general. We want to describe this operator locally. Following the analysis in [Mül11], we consider an open subset U of X such that $E_\chi|_U$ is trivial. Then, $E_0 \otimes E_\chi|_U$ is isomorphic to the direct sum of m -copies of $E_0|_U$, i.e.,

$$(E_0 \otimes E_\chi)|_U \cong \oplus_{i=1}^m E_0|_U,$$

where $m := \text{rank}(E_\chi) = \dim V_\chi$. Let $(e_i), i = 1, \dots, m$ be any basis of flat sections of $E_\chi|_U$. Then, each $\phi \in C^\infty(U, (E_0 \otimes E_\chi)|_U)$ can be written as

$$\phi = \sum_{i=1}^m \phi_i \otimes e_i,$$

where $\phi_i \in C^\infty(U, E_0|_U), i = 1, \dots, m$. The product connection is given by

$$\nabla_Y^{E_0 \otimes E_\chi} \phi = \sum_{i=1}^m (\nabla_Y^{E_0})(\phi_i) \otimes e_i,$$

where $Y \in C^\infty(X, TX)$. The local expression above is independent of the choice of the base of flat sections of $E_\chi|_U$, since the transition maps comparing flat sections are constant. By (4.1), we obtain the twisted Bochner-Laplace operator acting on $C^\infty(X, E_0 \otimes E_\chi)$ given by

$$\Delta_{E_0, \chi}^\sharp \phi = \sum_{i=1}^m (\Delta_{E_0} \phi_i) \otimes e_i, \quad (4.2)$$

where Δ_{E_0} denotes the Bochner-Laplace operator $\Delta^{E_0} = (\nabla^{E_0})^* \nabla^{E_0}$ associated to the connection ∇^{E_0} . Let now $\tilde{E}_0, \tilde{E}_\chi$ be the pullbacks to \tilde{X} of E_0, E_χ , respectively. Then,

$$\tilde{E}_\chi \cong \tilde{X} \times V_\chi,$$

and

$$C^\infty(\tilde{X}, \tilde{E}_0 \otimes \tilde{E}_\chi) \cong C^\infty(\tilde{X}, \tilde{E}_0) \otimes V_\chi. \quad (4.3)$$

With respect to the isomorphism (4.3), it follows from (4.2) that the lift of $\Delta_{E_0, \chi}^\sharp$ to \tilde{X} takes the form

$$\tilde{\Delta}_{E_0, \chi}^\sharp = \tilde{\Delta}_{E_0} \otimes \text{Id}_{V_\chi}, \quad (4.4)$$

where $\tilde{\Delta}_{E_0}$ is the lift of Δ_{E_0} to \tilde{X} . By (4.2), $\Delta_{E_0, \chi}^\sharp$ has principal symbol

$$\sigma_{\Delta_{E_0, \chi}^\sharp}(x, \xi) = \|\xi\|_x^2 \text{Id}_{(E_0 \otimes E_\chi)_x}, \quad x \in X, \xi \in \mathbb{T}_x^* X.$$

Hence, it has nice spectral properties, i.e., its spectrum is discrete and contained in a translate of a positive cone $C \subset \mathbb{C}$ such that $\mathbb{R}^+ \subset C$ ([Spi18, Lemma 8.6]).

We specialize now the twisted Bochner-Laplace operator $\Delta_{E_0, \chi}^\sharp$ to the case of the operator $\Delta_{\tau, \chi}^\sharp$ acting on smooth sections of the twisted vector bundle $E_\tau \otimes E_\chi$. Here, E_τ is the locally homogeneous vector bundle associated with a finite-dimensional, unitary representation τ of K . The keypoint is that when we consider the lift $\tilde{\Delta}_{\tau, \chi}^\sharp$ of the twisted Bochner-Laplace operator $\Delta_{\tau, \chi}^\sharp$ to the universal covering \tilde{X} , it acts as the identity operator on V_χ . By (4.1), we have

$$\tilde{\Delta}_{\tau, \chi}^\sharp = \tilde{\Delta}_\tau \otimes \text{Id}_{V_\chi},$$

where $\tilde{\Delta}_\tau$ is the lift to \tilde{X} of the Bochner-Laplace operator Δ_τ , associated with the representation τ of K .

4.2 The twisted operator $A_\chi^\sharp(\sigma)$

In this section, we define the twisted operators $A_\chi^\sharp(\sigma)$, associated with $\sigma \in \widehat{M}$ and representations χ of Γ , acting on smooth sections of twisted vector bundles. These operators are first introduced in [BO95]. For more details we refer the reader to [Spi18, Section 5].

We define the restricted Weyl group as the quotient $W_A := M'/M$. Then, W_A has order 2. Let $w \in W_A$ be the non-trivial element of W_A , and m_w a representative of w in M' . The action of W_A on \widehat{M} is defined by $(w\sigma)(m) := \sigma(m_w^{-1}mm_w)$, $m \in M, \sigma \in \widehat{M}$. Following the proof of Proposition 1.1 in [BO95] (see also [Pfa12, Proposition 2.3]), there exist unique integers $m_\tau(\sigma) \in \{-1, 0, 1\}$, which are equal to zero except for finitely many $\tau \in \widehat{K}$, such that there exist unique integers $m_\tau(\sigma) \in \{-1, 0, 1\}$, which are equal to zero except for finitely many $\tau \in \widehat{K}$, such that,

- if σ is Weyl invariant, $\sigma = \sum_{\tau \in \widehat{K}} m_\tau(\sigma) i^*(\tau)$;
- if σ is non-Weyl invariant, $\sigma + w\sigma = \sum_{\tau \in \widehat{K}} m_\tau(\sigma) i^*(\tau)$.

We define a locally homogeneous vector bundle $E(\sigma)$ associated to σ by

$$E(\sigma) := \bigoplus_{\substack{\tau \in \widehat{K} \\ m_\tau(\sigma) \neq 0}} E_\tau,$$

where E_τ is the locally homogeneous vector bundle associated with $\tau \in \widehat{K}$. The vector bundle $E(\sigma)$ has a grading

$$E(\sigma) = E(\sigma)^+ \oplus E(\sigma)^-. \quad (4.5)$$

This grading is defined exactly by the positive or negative sign of $m_\tau(\sigma)$. Let $\tilde{E}(\sigma)$ be the pullback of $E(\sigma)$ to \tilde{X} . Then,

$$\tilde{E}(\sigma) = \bigoplus_{\substack{\tau \in \widehat{K} \\ m_\tau(\sigma) \neq 0}} \tilde{E}_\tau.$$

We assume now that $\tau \in \widehat{K}$ is irreducible. Recall that $\tilde{\Delta}_\tau = -R(\Omega) + \lambda_\tau \text{Id}$. ([Mia80, Proposition 1.1], see also [Spi18, (5.4)]) We put

$$\tilde{A}_\tau := \tilde{\Delta}_\tau - \lambda_\tau \text{Id}.$$

Hence, the operator \tilde{A}_τ acts like $-R(\Omega)$ on the space of smooth sections of \tilde{E}_τ . It is an elliptic, formally self-adjoint operator of second order. By [Che73], it is an essentially self-adjoint operator. Its self-adjoint extension will be also denoted by \tilde{A}_τ . We get then the operator $\tilde{A}_{\tau,\chi}^\sharp$ acting on the space $C^\infty(\tilde{X}, \tilde{E}_\tau \otimes \tilde{E}_\chi)$, defined by

$$\tilde{A}_{\tau,\chi}^\sharp = \tilde{A}_\tau \otimes \text{Id}_{V_\chi}.$$

We put

$$c(\sigma) := -|\rho|^2 - |\rho_m|^2 + |\nu_\sigma + \rho_m|^2, \quad (4.6)$$

where ν_σ is as in (2.5) and ρ, ρ_m are defined by (2.3) and (2.4), respectively. We define the operator $A_\chi^\sharp(\sigma)$ acting on $C^\infty(X, E(\sigma) \otimes E_\chi)$ by

$$A_\chi^\sharp(\sigma) := \bigoplus_{m_\tau(\sigma) \neq 0} A_{\tau,\chi}^\sharp + c(\sigma). \quad (4.7)$$

The operator $A_\chi^\sharp(\sigma)$ preserves the grading and it is a non-self-adjoint, elliptic operator of order two.

5 The determinant formula

In [Spi15], a determinant formula has been proved, which gives an interpretation of the twisted Ruelle zeta function in terms of graded, regularized determinants of twisted, non-self-adjoint, elliptic differential operators. We recall here some facts and definitions from [Spi15].

Let σ_p be the standard representation of M in $\Lambda^p \mathbb{R}^{d-1} \otimes \mathbb{C}$. Let $\sigma, \sigma' \in \widehat{M}$. We denote by $[\sigma_p \otimes \sigma : \sigma']$ the multiplicity of σ' in $\sigma_p \otimes \sigma$. We distinguish again two cases for $\sigma' \in \widehat{M}$.

- **case (a):** σ' is invariant under the action of the restricted Weyl group W_A . Then, $i^*(\tau) = \sigma'$, where $\tau \in R(K)$.
- **case (b):** σ' is not invariant under the action of the restricted Weyl group W_A . Then, $i^*(\tau) = \sigma' + w\sigma'$, where $\tau \in R(K)$.

We define the operator

$$A_\chi^\sharp(\sigma_p \otimes \sigma) := \bigoplus_{[\sigma'] \in \widehat{M}/W_A} \bigoplus_{i=1}^{[\sigma_p \otimes \sigma : \sigma']} A_\chi^\sharp(\sigma'), \quad (5.1)$$

acting on the space $C^\infty(X, E(\sigma') \otimes E_\chi)$, where $E(\sigma')$ is the vector bundle over X , constructed as in [Spi18, p. 175].

Let α be the unique positive root of $(\mathfrak{g}, \mathfrak{a})$. Let $H \in \mathfrak{a}$ such that $\alpha(H) = 1$. The character $\lambda \equiv \lambda_p$ of A is defined by $\lambda \equiv \lambda_p(a) = e^{p\alpha(\log a)}$. Then, we can identify λ with p . By [Spi15, Proposition 7.9, case (a)], we have the following proposition.

Proposition 5.1. *The Ruelle zeta function has the representation*

$$R(s; \sigma, \chi) = \prod_{p=0}^{d-1} \det_{\text{gr}}(A_\chi^\sharp(\sigma_p \otimes \sigma) + (s + |\rho| - p)^2)^{(-1)^p} \cdot \exp\left(\left(-1\right)^{\frac{d-1}{2}+1} \pi(d+1) \dim(V_\sigma) \dim(V_\chi) \frac{\text{Vol}(X)}{\text{Vol}(S^d)} s\right). \quad (5.2)$$

Remark 5.2. In the determinant formula (5.1), the regularized determinants are graded, with respect to the grading (4.5) of the locally homogeneous vector bundle $E(\sigma)$ over X .

Theorem 5.3. *The Ruelle zeta function has the representation*

$$R(s; \chi) = \prod_{k=0}^{d-1} \prod_{p=k}^{d-1} \det \left(\Delta_{\chi, k}^{\sharp} + s(s + 2(|\rho| - p)) \right)^{(-1)^p} \cdot \exp \left((-1)^{\frac{d-1}{2}+1} \pi(d+1) \dim(V_{\chi}) \frac{\text{Vol}(X)}{\text{Vol}(S^d)} s \right). \quad (5.3)$$

Let $d_{\chi, k} := \dim \text{Ker}(\Delta_{\chi, k})$. Then, the singularity of the Ruelle zeta function at $s = 0$ is of order

$$\sum_{k=0}^{(d-1)/2} (d+1-2k)(-1)^k d_{\chi, k}. \quad (5.4)$$

Proof. We denote by r_p be the p -th exterior power of the standard representation of K . This notation coincides with the notation used in [BO95, p. 23–24]. Then, for $p = 0, \dots, d-1$, we have $i^*(r_p) = \sigma_p + \sigma_{p-1}$. Let $k_p \in R(K)$, defined by

$$k_p := \sum_{k=0}^p (-1)^k r_{p-k}.$$

Then, for $p = 0, 1, \dots, d-1$, one has $\sigma_p = i^*(k_p)$. Let E_{k_p}, E_{r_k} be the locally homogeneous vector bundles over X , associated with k_p and r_k , correspondingly. We consider σ being trivial. By (4.6), $c(\sigma_p) = (|\rho| - p)^2$. By (4.7) and (5.1), the operators $A_{\chi}^{\sharp}(\sigma_p) + (|\rho| - p)^2$ and $-R(\Omega)$ on $C^{\infty}(X, E_{k_p} \otimes V_{\chi})$ coincide. On the other hand, by Kuga's Lemma (Subsection 2.1.3), the Casimir operator $R(\Omega)$ acts as the negative Hodge Laplacian $-\Delta_{\chi, k}^{\sharp}$ on $C^{\infty}(X, E_{r_k} \otimes V_{\chi})$. Hence, by (5.2), we get

$$\begin{aligned} R(s; \chi) &= \prod_{p=0}^{d-1} \det_{\text{gr}} \left(A_{\chi}^{\sharp}(\sigma_p) + (s + |\rho| - p)^2 \right)^{(-1)^p} \\ &\quad \cdot \exp \left((-1)^{\frac{d-1}{2}+1} \pi(d+1) \dim(V_{\chi}) \frac{\text{Vol}(X)}{\text{Vol}(S^d)} s \right) \\ &= \prod_{k=0}^{d-1} \prod_{p=k}^{d-1} \det \left(\Delta_{\chi, k}^{\sharp} + s(s + 2(|\rho| - p)) \right)^{(-1)^p} \\ &\quad \cdot \exp \left((-1)^{\frac{d-1}{2}+1} \pi(d+1) \dim(V_{\chi}) \frac{\text{Vol}(X)}{\text{Vol}(S^d)} s \right). \end{aligned}$$

For the order of the singularity of the Ruelle zeta function at $s = 0$, (5.4) follows from Poincaré duality on differential forms. \square

Remark 5.4. We note here that for $p = \frac{d-1}{2}$, σ_p is not irreducible. In fact, $\sigma_{\frac{d-1}{2}}$ decomposes into $\sigma_+ + \sigma_-$, and $\sigma_{\pm} = w\sigma_{\mp}$. Then, we use (5.1) to define $A_{\chi}^{\sharp}(\sigma_{\frac{d-1}{2}})$. By [Mil78, Lemma 1.4], $c(\sigma_+) = c(\sigma_-) = 0$.

Remark 5.5. In the case of unitary representations, which are acyclic, one can obtain the triviality of the kernels of the Hodge Laplacian $\Delta_{\chi,k}$, since by Hodge theorem

$$H^k(X, E_\chi) \cong \text{Ker}(\Delta_{\chi,k}) = \{0\},$$

where $H^k(X, E_\chi)$ is the cohomology with coefficients in the local system defined by χ . In the present case, there is no such an isomorphism and one can find non-injective Laplacians associated with acyclic representations of Γ .

6 Spaces of representations of the fundamental group

6.1 Odd signature operator

Following the idea of Braverman and Kappeler in [BK08b], we consider representations of Γ , which are non-unitary, but belong to a neighbourhood of the set of acyclic, unitary representations in a suitable topology, such that the odd signature operator B (see Definition 6.1.1 below) is bijective. For these representations, Braverman and Kappeler defined the refined analytic torsion for a compact, oriented, odd-dimensional manifold X . Having applications in mind, instead of considering the regularized determinant of the (flat) Laplacian they consider the graded determinant of the even part of its square, the odd signature operator B . We recall here some definitions from [BK08b] and introduce a more general geometrical setting.

Let X be a compact, oriented, Riemannian manifold of odd dimension $d = 2r - 1$. Denote by g the Riemannian metric on X . Let $E \rightarrow X$ be a complex vector bundle over X , endowed with a flat connection ∇ . Let $\Lambda^k(X, E)$ be the space of k -differential forms on X with values in E . We denote by ∇ the induced differential $\nabla: \Lambda^k(X, E) \rightarrow \Lambda^{k+1}(X, E)$. Let $\Gamma: \Lambda^k(X, E) \rightarrow \Lambda^{d-k}(X, E)$ be the chirality operator defined by the formula

$$\Gamma\omega := i^{(d+1)/2}(-1)^{k(k+1)/2} *' \omega,$$

where $*'$ denotes the operator acting on sections of $\Lambda^k T^*X \otimes E$ as $* \otimes \text{Id}$, and $*$ is the usual Hodge $*$ -operator.

Definition 6.1.1. We define the odd signature operator $B = B(\nabla, g)$ acting on $\Lambda^*(X, E)$ by

$$B := \Gamma\nabla + \nabla\Gamma. \tag{6.1}$$

We set

$$\Lambda^{\text{even}}(X, E) := \bigoplus_{p=0}^{r-1} \Lambda^{2p}(X, E).$$

Let B^{even} be the even part of the odd signature operator B , defined by $B^{\text{even}} := B: \Lambda^{\text{even}}(X, E) \rightarrow \Lambda^{\text{even}}(X, E)$.

6.2 Spaces of connections

We consider the following two assumptions.

Assumption 1. The connection ∇ is acyclic, i.e., the twisted de Rham complex

$$0 \rightarrow \Lambda^0(X, E) \xrightarrow{\nabla} \Lambda^1(X, E) \xrightarrow{\nabla} \dots \xrightarrow{\nabla} \Lambda^d(X, E) \rightarrow 0$$

is acyclic,

$$\text{Im}(\nabla|_{\Lambda^{k-1}(X, E)}) = \text{Ker}(\nabla|_{\Lambda^k(X, E)}),$$

for every $k = 1, \dots, d$.

Assumption 2. The odd signature operator $B: \Lambda^k(X, E) \rightarrow \Lambda^k(X, E)$ is bijective.

Suppose that there exists a Hermitian metric h on E , which is preserved by ∇ . In such a case, we call ∇ a Hermitian connection. For such connections, one can easily see that Assumption 1 implies Assumption 2 and vice versa (see [BK08b, Subsection 6.6]). Hence, all acyclic, Hermitian connections satisfy Assumptions 1 and 2.

Following [BK08b], we denote by $\Lambda^1(X, \text{End}(E))$ be the space of 1-forms with values in $\text{End}(E)$. We define the sup-norm on $\Lambda^1(X, \text{End}(E))$ by

$$\|\omega\|_{\text{sup}} := \max_{x \in X} |\omega(x)|,$$

where the norm $|\cdot|$ is induced by a Hermitian metric on E and a Riemannian metric on X . The topology defined by this norm is called the C^0 -topology and it is independent of the metrics. We identify the space of connections on E with $\Lambda^1(X, \text{End}(E))$, by choosing a connection ∇_0 and associating to a connection ∇ the 1-form $\nabla - \nabla_0 \in \Lambda^1(X, \text{End}(E))$. Hence, by this identification, the C^0 -topology on $\Lambda^1(X, \text{End}(E))$ provides a topology on the space of connections on E , which is independent of the choice of ∇_0 . This topology is called the C^0 -topology on the space of connections.

Let $\text{Flat}(E)$ be the set of flat connections on E and $\text{Flat}'(E, g) \subset \text{Flat}(E)$ be the set of flat connections on E , satisfying Assumptions 1 and 2. The topology induced on these sets by the C^0 -topology on the space of connections on E is also called the C^0 -topology. The following proposition is proved in [BK08b].

Proposition 6.2.1. *Flat'(E, g) is an C^0 -open subset of Flat(E), which contains all acyclic Hermitian connections on E.*

Proof. See [BK08b, Proposition 6.8]. □

6.3 Spaces of representations

Let $\text{Rep}(\pi_1(X), \mathbb{C}^n)$ be the space of all n -dimensional, complex representations of $\pi_1(X)$. This space has a natural structure of a complex algebraic variety. Let $\{\gamma_1, \dots, \gamma_L\}$ be a finite set of generators of $\pi_1(X)$. The elements γ_i satisfy finitely many relations. Then, a representation $\chi \in \text{Rep}(\pi_1(X), \mathbb{C}^n)$ is given by $2L$ invertible $n \times n$ -matrices $\chi(\gamma_1), \dots, \chi(\gamma_L), \chi(\gamma_1^{-1}), \dots, \chi(\gamma_L^{-1})$, which satisfy finitely many polynomial equations. Hence, we view $\text{Rep}(\pi_1(X), \mathbb{C}^n)$ as an algebraic subset of $\text{Mat}_{n \times n}(\mathbb{C})^{2L}$ with the induced topology.

For a representation $\chi \in \text{Rep}(\pi_1(X), \mathbb{C}^n)$, we consider the associated flat vector bundle $E_\chi \rightarrow X$. Let ∇_χ be the flat connection on E_χ . Then, the

monodromy of ∇_χ is isomorphic to χ . We denote also by ∇_χ the induced differential $\nabla_\chi: \Lambda^*(X, E_\chi) \rightarrow \Lambda^{*+1}(X, E_\chi)$. We denote by B_χ the odd signature operator $B_\chi = B(\nabla_\chi, g)$ acting on $\Lambda^*(X, E_\chi)$. Let B_χ^{even} be the restriction of B_χ to $\Lambda^{\text{even}}(X, E_\chi)$.

Let $\text{Rep}_0(\pi_1(X), \mathbb{C}^n)$ be the set of all acyclic representations of $\pi_1(X)$, i.e., the set of all representations χ such that ∇_χ is acyclic. Let $\text{Rep}^u(\pi_1(X), \mathbb{C}^n)$ be the set of all unitary representations of $\pi_1(X)$, i.e., the set of all representations χ such that there exists a Hermitian scalar product (\cdot, \cdot) on \mathbb{C}^n , which is preserved by the matrices $\chi(\gamma)$, for all $\gamma \in \pi_1(X)$. $\text{Rep}^u(\pi_1(X), \mathbb{C}^n)$ can be viewed as the real locus of the complex algebraic variety $\text{Rep}(\pi_1(X), \mathbb{C}^n)$. We set

$$\text{Rep}_0^u(\pi_1(X), \mathbb{C}^n) := \text{Rep}_0(\pi_1(X), \mathbb{C}^n) \cap \text{Rep}^u(\pi_1(X), \mathbb{C}^n).$$

Suppose that for some representation $\chi_0 \in \text{Rep}_0(\pi_1(X), \mathbb{C}^n)$ the operator B_{χ_0} is bijective. Then, there exists an open neighbourhood (in classical topology) $V \subset \text{Rep}(\pi_1(X), \mathbb{C}^n)$ of the set $\text{Rep}_0^u(\pi_1(X), \mathbb{C}^n)$ of acyclic, unitary representations such that, for all $\chi \in V$, the pair (∇_χ, g) satisfies Assumptions 1 and 2 of Subsection 6.2 (see [BK08b, Subsection 13.7]).

7 Refined analytic torsion

7.1 Definition

Let ∇ be acyclic. Let $\mathcal{M}(\nabla)$ be the set of all Riemannian metrics g on X , such that $B^{\text{even}} = B^{\text{even}}(\nabla, g)$ is bijective. Then, by [BK08b, Proposition 6.8], $\mathcal{M}(\nabla) \neq \emptyset$, for all flat connections in an open neighbourhood, in C^0 -topology, of the set of acyclic Hermitian connections. Let V be an open neighbourhood of $\text{Rep}_0^u(\pi_1(X), \mathbb{C}^n)$ as in Subsection 6.3.

Definition 7.1.1. Let $\chi \in V$. We define the refined analytic torsion $T_\chi = T(\nabla_\chi)$ by

$$T_\chi := \det_{\text{gr}, \theta}(B_\chi^{\text{even}}) e^{i\pi \text{rank}(E_\chi) \eta_{tr}} \in \mathbb{C} \setminus \{0\}, \quad (7.1)$$

where $\eta_{tr} = \eta_{tr}(g) = \frac{1}{2} \eta(0, B_{tr}(g))$, and $\eta(0, B_{tr}(g))$ denotes the eta invariant of the even part of the odd signature operator corresponding to the trivial line bundle, endowed with the trivial connection. The refined analytic torsion is independent of the Agmon angle $\theta \in (-\pi, 0)$ for the operator $B_\chi^{\text{even}} = B_\chi^{\text{even}}(g)$ and the Riemannian metric $g \in \mathcal{M}(\nabla)$.

Remark 7.1.2. The graded determinant of the operator B_χ^{even} is slightly different than the usual expression of the graded determinant as in Subsection 2.2.3. We recall here briefly the definition of $\det_{\text{gr}}(B_\chi^{\text{even}})$. For more details, we refer the reader to [BK08b, Subsection 2.2 and 6.11]. We set

$$\begin{aligned} \Lambda_+^k(X, E) &:= \text{Ker}(\nabla\Gamma) \cap \Lambda^k(X, E) \\ \Lambda_-^k(X, E) &:= \text{Ker}(\Gamma\nabla) \cap \Lambda^k(X, E). \end{aligned}$$

Assumption 2 implies $\Lambda^k(X, E) = \Lambda_+^k(X, E) \oplus \Lambda_-^k(X, E)$ (see [BK08b, Subsection 6.9]). We set

$$\Lambda_\pm^{\text{even}}(X, E) = \bigoplus_{p=0}^{r-1} \Lambda_\pm^{2p}(X, E).$$

We denote by $B_{\chi,\pm}^{\text{even}}$ the restriction of B_χ^{even} to $\Lambda_\pm^{\text{even}}(X, E)$. Then, B_χ^{even} leaves the subspaces $\Lambda_\pm^{\text{even}}(X, E)$ invariant. By Assumption 2, the operators $B_{\chi,\pm}^{\text{even}}$ are bijective. Hence, we define the graded determinant $\det_{\text{gr}}(B_\chi^{\text{even}}) \in \mathbb{C} \setminus \{0\}$ of B_χ^{even} by

$$\det_{\text{gr},\theta}(B_\chi^{\text{even}}) := \frac{\det_\theta(B_{\chi,+}^{\text{even}})}{\det_\theta(-B_{\chi,-}^{\text{even}})},$$

where θ is an Agmon angle for B_χ^{even} (and an Agmon angle for $B_{\chi,\pm}^{\text{even}}$ as well).

Remark 7.1.3. By [BK08b, Corollary 13.11.(2)], the refined analytic torsion T_χ is an holomorphic function on the set $V \setminus \Sigma$, where Σ denotes the set of singular points of the complex algebraic variety $\text{Rep}(\pi_1(X), \mathbb{C}^n)$.

7.2 Independence of the refined analytic torsion of the Riemannian metric and the Agmon angle

The independence of T_χ of the Agmon angle θ and the Riemannian metric g is proved in [BK08b, Theorem 9.3]. We consider the eta invariant $\eta = \eta(\chi, g) = \eta(B_\chi)$ of the even part of the odd signature operator. We set

$$\xi = \xi(\chi, g, \theta) := \frac{1}{2} \sum_{k=0}^d (-1)^{k+1} k L \det_{2\theta}(B_\chi^2|_{\Lambda^k(X, E_\chi)}). \quad (7.2)$$

The number ξ is defined in [BK08b, (7.79)] (see also (8.102) in the same paper). Let $\eta = \eta(\chi, g) = \eta(B_\chi^{\text{even}})$ denote the eta invariant of the even part of the odd signature operator B_χ . By [BK08b, Theorem 7.2], for a suitable choice of an Agmon angle for B_χ , the following equality holds

$$\det_{\text{gr},\theta}(B_\chi^{\text{even}}) = e^{\xi(\chi, g, \theta)} e^{-i\pi\eta(\chi, g)}. \quad (7.3)$$

The graded determinant $\det_{\text{gr}}(B_\chi^{\text{even}})$ above does depend on the Riemannian metric g . This is the reason why the additional exponential $e^{i\pi \text{rank}(E_\chi)\eta_{\text{tr}}(g)}$ is considered in the definition of the refined analytic torsion in [BK08b], to remove the metric anomaly. By results in [Gil84, p. 52], modulo \mathbb{Z} , the difference

$$\eta(\chi, g) - \text{rank}(E_\chi)\eta_{\text{tr}}(g)$$

is independent of g . In addition, by [BK08b, Proposition 9.7], for $g_1, g_2 \in \mathcal{M}(\nabla)$ and suitable choices of Agmon angles θ_0, θ_1 ,

$$\xi(\chi, g_1, \theta_1) = \xi(\chi, g_2, \theta_2) \pmod{\pi i}.$$

Hence, for different choices of Agmon angles and Riemannian metrics, the corresponding expressions in (7.1) coincide up to a sign. To see that the two expressions coincide see [BK08b, p. 238].

Remark 7.2.1.

$$\rho_\chi := \eta(\chi, g) - \text{rank}(E_\chi)\eta_{\text{tr}}(g),$$

is called the *rho* invariant ρ_χ of the operator B_χ^{even} .

8 Alternative definition of the refined analytic torsion

In [BK08b, Section 11], an alternative definition of the refined analytic torsion is given, which involves the Hirzebruch L -polynomial and the signature theorem by Atiyah, Patodi and Singer. We recall this definition here.

Let N be a smooth, oriented, compact, even-dimensional manifold such that $\partial N = X$. Recall that the signature $\text{sign}(N)$ is an integer defined in pure cohomological terms (see [APS75a, p. 65] and [APS75b, p. 407]), as the signature of the Hermitian form, induced by the cup product, in the middle $(\dim N/2)$ -cohomology, and it is metric independent. We denote by $L(p)$ the Hirzebruch L -polynomial in the Pontrjagin forms of a Riemannian metric of a manifold ([LM89, p. 228, 232]). The signature theorem ([APS75a, Theorem 4.14] , [APS75b, Theorem 2.2]) states

$$\text{sign}(N) = \int_N L(p) - \eta(B_{\text{trivial}}), \quad (8.1)$$

where $L_N(p) := L(p)$ denotes the Hirzebruch L -polynomial in the Pontrjagin forms of a Riemannian metric on N which is a product near X . By the metric independence of $\text{sign}(N)$, (8.1) and Proposition 9.5 in [BK08b], we have that modulo \mathbb{Z} ,

$$\eta - \text{rank}(E) \int_N L(p)$$

is independent of the Riemannian metric on X .

In general, there might be no smooth, oriented manifold N such that $\partial N = X$. However, since $\dim X$ is odd, there exists an oriented manifold N , whose oriented boundary is the disjoint union of two copies of X , with the same orientation ([Wal60], [RR98, Th. IV.6.5]). Then, as discussed above, modulo \mathbb{Z} ,

$$\eta - \frac{\text{rank}(E)}{2} \int_N L(p).$$

is independent of the Riemannian metric on X .

We assume that the flat connection ∇ on E belongs to an open neighbourhood, in C^0 -topology, of the set of acyclic Hermitian connections (see Subsection 6.2).

Definition 8.1. *Let $\theta \in (-\pi, 0)$ be the Agmon angle of B^{even} . Choose a smooth, compact, oriented manifold N , whose oriented boundary is diffeomorphic to two disjoint copies of M . The refined analytic torsion $T'(\nabla)$ is defined by*

$$T'(\nabla) = T'(X, E, \nabla, N) := \det_{\text{gr}, \theta}(B^{\text{even}}) e^{i\pi \frac{\text{rank}(E)}{2} \int_N L(p)}, \quad (8.2)$$

where where $L_N(p) := L(p)$ denotes the Hirzebruch L -polynomial in the Pontrjagin forms of a Riemannian metric on N which is a product near ∂N .

By the discussion above and the same arguments as in the proof of [BK08b, Theorem 9.3], $T'(\nabla)$ is independent the Agmon angle θ and of the Riemannian metric on X . However, $T'(\nabla)$ does depend on the choice of N . Since for different choices of N , $\int_N L(p)$ differs by an integer, $T'(\nabla)$ is independent of the choice of

N up to a multiplication by $i^k \text{rank}(E)$, $k \in \mathbb{Z}$. Hence, if $\text{rank}(E)$ is even, $T'(\nabla)$ is well defined up to a sign, and if $\text{rank}(E)$ is divisible by 4, it is a well defined complex number. In addition, by (8.2) and [BK08b, Theorem 7.2], we have

$$T'(\nabla) = e^\xi e^{-i\pi\eta} e^{i\pi \frac{\text{rank}(E)}{2}} \int_N L(p), \quad (8.3)$$

where ξ is as in (7.2).

9 Cappell-Miller complex torsion

In [CM10], Cappell and Miller introduced the Cappell-Miller torsion τ , an invariant, which is an element of $\det(H^*(X, E)) \otimes \det(H^*(X, E))$. To define this invariant, they used the flat Laplacian on vector bundle-valued differential forms. This operator is non-self-adjoint and in terms of the chirality operator in the previous sections is the square B^2 of the odd signature operator B . Hence, roughly speaking, Braverman and Kappeler used the odd signature operator to define the refined analytic torsion and Cappell and Miller its square. By [BK08a, eq. (5.1)], for a finite-dimensional complex (C^*, ∂) ,

$$\tau = \tau_\Gamma := \rho_\Gamma \otimes \rho_\Gamma \in \det(H^*(\partial)) \otimes \det(H^*(\partial)),$$

where $\rho_\Gamma \in \det(H^*(d))$ is as in Definition 11.1.1 (Appendix). If the complex (C^*, d) is acyclic then τ_Γ is a complex number and by [BK08a, eq. (5.3)],

$$\tau_\Gamma := \prod_{k=0}^d \det(B^2|_{C^k})^{k(-1)^{k+1}} \quad (9.1)$$

Remark 9.1. As explained in [BK08a, Remark 5.2], the element τ in [CM10] is constructed slightly different. In particular, by the construction of the element ρ_Γ and the construction in [CM10, Section 6], the two elements coincide up to a sign. One can see that the signs agree by (9.1) and equation (6.18) in [CM10].

In the manifold setting, we consider the Cappell-Miller torsion $\tau_{\Gamma_{[0,\lambda]}} \in \det(H_{[0,\lambda]}^*(X, E)) \otimes \det(H_{[0,\lambda]}^*(X, E))$ of the finite-dimensional complex $\Lambda_{[0,\lambda]}^k(X, E)$ (see Appendix, Subsection 11.2).

Definition 9.2. Let $\theta \in (0, 2\pi)$ be an Agmon angle for the operator $B_{(\lambda, \infty)}^2$. Then, the Cappell-Miller torsion τ_∇ is defined by

$$\tau_\nabla := \tau_{\Gamma_{[0,\lambda]}} \prod_{k=0}^d \det_\theta(B^2|_{\Lambda_{(\lambda, \infty)}^k(X, E)})^{k(-1)^{k+1}} \in \det(H^*(X, E)) \otimes \det(H^*(X, E)).$$

By [CM10, Theorem 8.3], the Cappell-Miller torsion τ_∇ is independent of the choice of $\lambda > 0$ and of the choice of the Riemannian metric on X . Moreover, since the principal symbol of B^2 is self-adjoint, by [BK08b, Section 3.10], the regularized determinant $\det_\theta(B^2|_{\Lambda_{(\lambda, \infty)}^k(X, E)})$ is independent of the choice of the Agmon angle θ .

Remark 9.3. In the presence of the chirality operator Γ , the dual differential $\partial^{*,\sharp}$, introduced in [CM10] is given by $\partial^{*,\sharp} = \Gamma \partial \Gamma$. Then, $(\partial^{*,\sharp})^2 = 0$ and one

obtains a complex $(\Lambda^*(X, E), \partial^{*,\sharp})$ of degree -1 . In addition, the flat Laplacian Δ^\sharp acting on $\Lambda^*(X, E)$ is given by

$$\Delta^\sharp := \partial\partial^{*,\sharp} + \partial^{*,\sharp}\partial.$$

This operator is the same as the square B^2 of the odd signature operator, introduced in Section 6. Then, the Cappell-Miller torsion τ is defined in terms of the bicomplex $(\Lambda^*(X, E), \partial, \partial^{*,\sharp})$. In particular, it is given as a combination of the **torsion** $(\Lambda_{[0,\lambda]}^k(X, E), \partial, \partial^{*,\sharp}) \in \det(H^*(X, E)) \otimes \det(H^*(X, E))$ and the square of the **Ray – Singer** term, which corresponds to the terms $\tau_{\Gamma_{[0,\lambda]}}$ and $\prod_{k=0}^d \det\theta(B^2|_{\Lambda_{(\lambda,\infty)}^k(X, E)})^{k(-1)^{k+1}}$, respectively, in Definition 9.2.

We turn now to the case, where the flat connection ∇ on the flat vector bundle belongs to the neighbourhood of the set of all acyclic Hermitian connections. Let $\chi \in V$, where V is as in Subsection 6.3. Then, B_χ is bijective, the cohomologies $H^*(X, E_\chi)$ vanish and the element $\tau_{\Gamma_{[0,\lambda]}}$ is equal to 1. For $\chi \in V$, we define the Cappell-Miller torsion $\tau_\chi := \tau(\nabla_\chi)$ by

$$\tau_\chi := \prod_{k=0}^d \det(B_{\chi,k}^2)^{k(-1)^{k+1}}, \quad (9.2)$$

where $B_{\chi,k}^2$ denotes the operator B_χ^2 acting on $\Lambda^k(X, E)$. In this case τ_χ is a complex number.

Proposition 9.4. *Let $\chi \in V$. Then, the Cappell-Miller torsion τ_χ and the refined analytic torsion T_χ are related by*

$$\tau_\chi = T_\chi^2 e^{2\pi i(\eta(B_\chi^{\text{even}}) - \text{rank}(E_\chi)\eta_{\text{tr}})}. \quad (9.3)$$

Proof. (9.3) follows from (7.1), (7.2), (7.3) and (9.2). \square

10 Ruelle zeta function and refined analytic torsion

Let now $X = \Gamma \backslash G/K$ be a compact, oriented, hyperbolic manifold, where G , K and Γ are as in Subsection 2.1.1. Let V be as in in Subsection 6.3, i.e., V is an open neighbourhood of the set $\text{Rep}_0^u(\pi_1(X), \mathbb{C}^n)$ of acyclic and unitary representations of $\pi_1(X) = \Gamma$, such that, for all $\chi \in V$, B_χ is bijective. Then, we have the following theorem.

Theorem 10.1. *Let $\chi \in V$. Then, the Ruelle zeta function $R(s; \chi)$ is regular at $s = 0$ and is equal to the complex Cappell-Miller torsion,*

$$R(0; \chi) = \tau_\chi. \quad (10.1)$$

Proof. For $\chi \in V$, the operator $B_{\chi,k}^2$ coincides with the flat Laplacian $\Delta_{\chi,k}^\sharp$ on $\Lambda^k(X, E_\chi)$ and it is injective. Hence, by Theorem 5.3, the Ruelle zeta function $R(s; \chi)$ is regular at zero and in addition, by (5.3),

$$R(0; \chi) = \prod_{k=0}^{d-1} \det(\Delta_{\chi,k}^\sharp)^{(d-k)(-1)^k} = \prod_{k=1}^d \det(\Delta_{\chi,k}^\sharp)^{k(-1)^{k-1}}. \quad (10.2)$$

Hence, (10.1) follows by (9.2) and (10.2). \square

Corollary 10.2. *Let $\chi \in V$. Then the Ruelle zeta function $R(s; \chi)$ is regular at $s = 0$ and is related to the refined analytic torsion T_χ by*

$$R(0; \chi) = T_\chi^2 e^{2\pi i(\eta(B_\chi^{\text{even}}) - \text{rank}(E_\chi)\eta_{\text{tr}})}. \quad (10.3)$$

Proof. The assertion follows from (10.1) and (9.3). \square

Let now $\chi \in \text{Rep}_0^u(\pi_1(X), \mathbb{C}^n)$. Then, by [BO95, Theorem 4.8], the Ruelle zeta function $R(s, \chi)$ associated with χ is regular at $s = 0$. Moreover, the Cappell-Miller torsion coincides with the square of the Ray-Singer real-valued torsion T_χ^{RS} . On the other hand, in this case, the refined analytic torsion is still a complex number, given by

$$T_\chi = e^{\xi_\chi} e^{-i\pi(\eta(B_\chi^{\text{even}}) - \text{rank}(E_\chi)\eta_{\text{tr}})}.$$

Here, we denote $\xi_\chi = \xi(\chi, g, \theta)$. If χ is unitary and acyclic, the term e^{ξ_χ} coincides with the Ray-Singer torsion T_χ^{RS} , and the eta invariant $\eta(\chi, g)$ is real valued. Hence,

$$|T_\chi| = e^{\xi_\chi} = T_\chi^{RS} \quad (10.4)$$

Therefore, (10.1), (10.3) and (10.4), yields the following classical result (see [BO95, Theorem 4.8]).

Corollary 10.3. *Let $\chi \in \text{Rep}_0^u(\pi_1(X), \mathbb{C}^n)$. Then, the Ruelle zeta function $R(s; \chi)$ associated with χ is regular at $s = 0$ and*

$$R(0; \chi) = \tau_\chi = |T_\chi|^2 = (T_\chi^{RS})^2.$$

Corollary 10.4. *Let $\chi \in V$. Then, the Ruelle zeta function $R(s; \chi)$ is regular at $s = 0$ and*

$$|R(0; \chi)| = |\tau_\chi| = (T_\chi^{RS})^2 \quad (10.5)$$

Proof. By [BK08b, Theorem 8.2], we have

$$e^{\text{Re}(\xi_\chi)} = T_\chi^{RS}. \quad (10.6)$$

Hence, (10.5) follows from (10.1), (10.3) and (10.6). \square

Using the alternative definition of the refined analytic torsion as in Section 8, we obtain the following corollary.

Corollary 10.5. *Let $\chi \in V$ and $n = 4k$, $k \in \mathbb{Z}$. Then, the Ruelle zeta function $R(s; \chi)$ is regular at $s = 0$ and*

$$R(0; \chi) = T_\chi'^2 e^{2\pi i(\eta(B_\chi^{\text{even}}) - \frac{\text{rank}(E_\chi)}{2} \int_N L(p))} \quad (10.7)$$

Proof. It follows from (10.2), the definition of ξ in (7.2) and the expression of $T'(\nabla)$ in (8.3). Here, we denote $T_\chi' = T'(\nabla_\chi)$. \square

11 Appendix: Refined analytic torsion as an element of the determinant line

For a general, non-unitary representation the definition of the refined analytic torsion differs from Definition 7.1 and it is not a complex number any more. It is an element of the determinant line of the cohomology. We recall here the definition from [BK07]. Let \mathbf{k} be a field of characteristic zero. Let V be a \mathbf{k} -vector space of dimension n . We define the determinant line of V by $\det(V^*) := \Lambda^n V$, where $\Lambda^n V$ denotes the n -th exterior power of V . We set $\det(0) := k$. If L is a \mathbf{k} -line (one-dimensional vector space), we define the dual line L^{-1} by $L^{-1} := \text{Hom}_{\mathbf{k}}(L, \mathbf{k})$. For $l \in L$, we denote by $l^{-1} \in L^{-1}$ the unique \mathbf{k} -linear map $L \rightarrow \mathbf{k}$, such that $l^{-1}(l) = 1$. For a graded \mathbf{k} -vector space $V^* = V^0 \oplus V^1 \dots \oplus V^d$, we define

$$\det(V^*) := \bigotimes_{j=0}^d \det(V^j)^{(-1)^j}.$$

Let (C^*, ∂) ,

$$0 \rightarrow C^0 \xrightarrow{\partial} C^1 \xrightarrow{\partial} \dots \xrightarrow{\partial} C^d \rightarrow 0,$$

be a complex of finite-dimensional \mathbf{k} -vector spaces. We call the integer d the length of the complex (C^*, ∂) . We denote by

$$H^*(\partial) = \bigoplus_{i=0}^d H^i(\partial)$$

the cohomology of (C^*, ∂) . We set

$$\begin{aligned} \det(C^*) &:= \bigotimes_{j=0}^d \det(C^j)^{(-1)^j} \\ \det(H^*) &:= \bigotimes_{j=0}^d \det(H^j(\partial))^{(-1)^j}. \end{aligned}$$

We want to consider the determinant line of a direct sum of two (or more) finite-dimensional \mathbf{k} -vector spaces. Let V, W be finite-dimensional \mathbf{k} -vector spaces of dimension $\dim V = k$ and $\dim W = l$, correspondingly. We define the canonical fusion isomorphism

$$\mu_{V,W}: \det(V) \otimes \det(W) \rightarrow \det(V \oplus W)$$

by

$$\begin{aligned} \mu_{V,W}: (v_1 \wedge v_2 \wedge \dots \wedge v_k) \otimes (w_1 \wedge w_2 \wedge \dots \wedge w_l) \\ \mapsto v_1 \wedge v_2 \wedge \dots \wedge v_k \wedge w_1 \wedge w_2 \wedge \dots \wedge w_l, \end{aligned}$$

where $v_i \in V$ and $w_j \in W$. For $v \in \det(V)$ and $w \in \det(W)$, we have

$$\mu_{V,W}(v \otimes w) = (-1)^{\dim V \dim W} \mu_{V,W}(w \otimes v).$$

We denote by

$$\mu_{V,W}^{-1}: \det(V)^{-1} \otimes \det(W)^{-1} \rightarrow \det(V \oplus W)^{-1}$$

the transpose of the inverse of $\mu_{V,W}$. Then, it follows that for $v \in \det(V)$ and $w \in \det(W)$

$$\mu_{V,W}^{-1}(v^{-1} \otimes w^{-1}) = (\mu_{V,W}(v \otimes w))^{-1}.$$

Let now V_1, \dots, V_r , finite-dimensional \mathbf{k} -vector spaces. We define an isomorphism

$$\mu_{V_1, \dots, V_r}: \det(V_1) \otimes \dots \otimes \det(V_r) \rightarrow \det(V_1 \oplus \dots \oplus V_r).$$

For $j = 1, \dots, r-1$, it holds

$$\mu_{V_1, \dots, V_r} = \mu_{V_1, \dots, V_{j-1}, V_j \oplus V_{j+1}, V_{j+2}, \dots, V_r} \circ (1 \otimes \dots \otimes 1 \otimes \mu_{V_j, V_{j+1}} \otimes 1 \otimes 1).$$

We fix now a direct sum decomposition

$$C^j = B^j \oplus H^j \oplus A^j, \quad j = 0, \dots, d,$$

such that $B^j \oplus H^j = \text{Ker}(\partial) \cap C^j$ and $B^j = \partial(C^{j-1}) = \partial(A^{j-1})$, for all j . Note that $A^d = 0$. Set $A^{-1} = \{0\}$. Then, H^j is naturally isomorphic to the cohomology $H^j(\partial)$ and ∂ defines an isomorphism $\partial: A^j \rightarrow B^{j+1}$. For each $j = 0, \dots, d$, we fix $c_j \in \det(C^j)$ and $a_j \in \det(A^j)$. Let $\partial(a_j) \in \det(B^{j+1})$ be the image of a_j under the map $\det(A^j) \rightarrow \det(B^{j+1})$, induced by the isomorphism $\partial: A^j \rightarrow B^{j+1}$. Then, for each $j = 0, \dots, d$, there is a unique element $h_j \in \det(H^j)$ such that

$$c_j = \mu_{B^j, H^j, A^j}(\partial(a_{j-1}) \otimes h_j \otimes a_j).$$

Let ϕ_{C^*} be the isomorphism

$$\phi_{C^*} = \phi_{(C^*, \partial)}: \det(C^*) \rightarrow \det(H^*(\partial)), \quad (11.1)$$

defined by

$$\phi_{C^*}: c_0 \otimes c_1^{-1} \otimes \dots \otimes c_d^{(-1)^d} \mapsto (-1)^{\mathcal{N}(C^*)} h_0 \otimes h_1^{-1} \otimes \dots \otimes h_d^{(-1)^d},$$

where

$$\mathcal{N}(C^*) := \frac{1}{2} \sum_{j=0}^d \dim A^j (\dim A^j + (-1)^{j+1}).$$

Then, ϕ_{C^*} is independent of the choices of c_j and a_j .

Remark 11.1. The isomorphism ϕ_{C^*} is a sign refinement of the standard construction by Milnor ([Mil66]). The sign factor $\mathcal{N}(C^*)$ was introduced by Braverman and Kappeler in order to obtain various compatibility properties (see [BK07, Remark 2.5, Lemma 2.7 and Proposition 5.6]).

11.1 Refined torsion of a finite-dimensional complex with a chirality operator

We consider now the chirality operator $\Gamma: C^* \rightarrow C^*$, such that $\Gamma: C^j \rightarrow C^{d-j}$, for $j = 0, \dots, d$. For $c_j \in \det(C^j)$, $\Gamma c_j \in \det(C^{d-j})$ is the image of c_j under the isomorphism $\det(C^j) \rightarrow \det(C^{d-j})$, induced by Γ . Let $d = 2r - 1$ be an odd

integer. We fix non-zero elements $c_j \in \det(C^j)$, $j = 0, \dots, r-1$, and consider the element

$$c_\Gamma := (-1)^{\mathcal{R}(C^*)} c_0 \otimes c_1^{-1} \otimes \dots \otimes c_{r-1}^{(-1)^{r-1}} \\ \otimes (\Gamma c_{r-1})^{(-1)^r} \otimes (\Gamma c_{r-2})^{(-1)^{r-1}} \otimes \dots \otimes (\Gamma c_0^{-1})$$

of $\det(C^*)$, where

$$\mathcal{R}(C^*) = \frac{1}{2} \sum_{j=0}^{r-1} \dim C^j (\dim C^j + (-1)^{r+j}).$$

It follows from the definition of c_j^{-1} that c_Γ is independent of the choice of c_j , $j = 0, \dots, r-1$.

Definition 11.1.1. We define the refined torsion ρ_Γ of the pair (C^*, Γ) by

$$\rho_\Gamma = \rho_{C^*, \Gamma} := \phi_{C^*}(c_\Gamma), \quad (11.2)$$

where ϕ_{C^*} is as in (11.1).

We consider now the finite-dimensional analogue of the odd signature operator B , defined in a Riemannian manifold-setting by (6.1). Let $B: C^* \rightarrow C^*$ be the signature operator, defined by

$$B := \partial\Gamma + \Gamma\partial.$$

We consider its square, $B^2 = (\partial\Gamma)^2 + (\Gamma\partial)^2$. Let $I \subset [0, \infty)$. Let $C_I^j \subset C^j$ be the span of the generalized eigenvectors of the restriction of B^2 to C^j , corresponding to eigenvalues r with $|r| \in I$. Both Γ and ∂ commute with B and hence with B^2 . Then, we have $\Gamma: C_I^j \rightarrow C_I^{d-j}$, and $\partial: C_I^j \rightarrow C_I^{j+1}$. Hence, we obtain a subcomplex C_I^* of C^* and the restriction Γ_I of Γ to C_I^* is the chirality operator on this complex. Let $\partial_I, B_I, B_I^{\text{even}}$ be the restriction to C_I^* of $\partial, B, B^{\text{even}}$, correspondingly. Here, we denote $B^{\text{even}}: C^{\text{even}} \rightarrow C^{\text{even}}$, where $C^{\text{even}} := \bigoplus_{j \text{ even}} C^j$. By Lemma 5.8 in [BK07], if $0 \notin I$, then the complex (C_I^*, ∂_I) is acyclic. For every $\lambda \geq 0$, we have

$$C^* = C_{[0, \lambda]}^* \oplus C_{(\lambda, \infty)}^*.$$

In particular,

$$H_{(\lambda, \infty)}^*(\partial) = 0, \quad H_{[0, \lambda]}^*(\partial) \cong H^*(\partial).$$

Hence, there are canonical isomorphisms

$$\Phi: \det(H_{(\lambda, \infty)}^*(\partial)) \rightarrow \mathbb{C}, \quad \Psi: \det(H_{[0, \lambda]}^*(\partial)) \rightarrow \det(H^*(\partial)).$$

By [BK07, Proposition 5.10], for each $\lambda \geq 0$,

$$\rho_\Gamma = \det_{\text{gr}}(B_{(\lambda, \infty)}^{\text{even}}) \rho_{\Gamma_{[0, \lambda]}},$$

where $\rho_{\Gamma_{[0, \lambda]}}$ is an element of $\det(H^*(\partial))$ via the canonical isomorphism Ψ .

11.2 Refined analytic torsion as an element of the determinant line for compact odd-dimensional manifolds

As in Section 6, we consider a compact, oriented, Riemannian manifold (X, g) of odd dimension $d = 2r - 1$ and a complex vector bundle E over X , endowed with a flat connection ∇ . Let $\Lambda^k(X, E)$ be the space of k -differential forms on X with values in E . Let B be the odd signature operator acting on $\Lambda^k(X, E)$ as in Definition 6.1. Let $I \subset [0, \infty)$. Let $\Lambda_I^k(X, E)$ be the image of $\Lambda^k(X, E)$ under the spectral projection of B^2 , corresponding to eigenvalues whose absolute value lie in I . If I is bounded, then the subspace $\Lambda_I^k(X, E)$ is finite-dimensional ([BK07, Section 6.10]). Let B_I, Γ_I be the restrictions to $\Lambda_I^k(X, E)$ of B and Γ , respectively. For every $\lambda \geq 0$, we have

$$\Lambda^*(X, E) = \Lambda_{[0, \lambda]}^*(X, E) \oplus \Lambda_{(\lambda, \infty)}^*(X, E).$$

Since for every $\lambda \geq 0$ the complex $\Lambda_{(\lambda, \infty)}^*(X, E)$ is acyclic, we get

$$H_{[0, \lambda]}^*(X, E) \cong H^*(X, E).$$

Definition 11.2.1. *The refined analytic torsion $T = T(\nabla)$ is an element of $\det(H^*(X, E))$, defined by*

$$T(\nabla) = \rho_{\Gamma_{[0, \lambda]}} \det_{\text{gr}}(B_{(\lambda, \infty)}^{\text{even}}) e^{i\pi \text{rank}(E_X) \eta_{\text{tr}}(g)},$$

where $\rho_{\Gamma_{[0, \lambda]}}$ is as in (11.2).

By [BK07, Proposition 7.8] and [BK07, Theorem 9.6], the refined analytic torsion is independent the choice of $\lambda \geq 0$, the choice of the Agmon angle $\theta \in (-\pi, 0)$ for B^{even} , and the Riemannian metric g .

References

- [APS75a] M. F. Atiyah, V. K. Patodi, and I. M. Singer, *Spectral asymmetry and Riemannian geometry. I*, Math. Proc. Cambridge Philos. Soc. **77** (1975), 43–69.
- [APS75b] Michael F Atiyah, Vijay Kumar Patodi, and IM Singer, *Spectral asymmetry and Riemannian geometry. II*, Mathematical Proceedings of the Cambridge Philosophical Society, vol. 78, Cambridge University Press, 1975, pp. 405–432.
- [BK07] Maxim Braverman and Thomas Kappeler, *Refined analytic torsion as an element of the determinant line*, Geom. Topol. **11** (2007), 139–213.
- [BK08a] ———, *A canonical quadratic form on the determinant line of a flat vector bundle*, International Mathematics Research Notices **2008** (2008), no. 9, rnn030–rnn030.
- [BK08b] ———, *Refined analytic torsion*, J. Differential Geom. **78** (2008), no. 2, 193–267.

- [BO95] Ulrich Bunke and Martin Olbrich, *Selberg zeta and theta functions*, Mathematical Research, vol. 83, Akademie-Verlag, Berlin, 1995, A differential operator approach.
- [CD19] Yann Chaubet and Nguyen Viet Dang, *Dynamical torsion for contact anosov flows*, arXiv preprint arXiv:1911.09931 (2019).
- [Che73] Paul R. Chernoff, *Essential self-adjointness of powers of generators of hyperbolic equations*, J. Functional Analysis **12** (1973), 401–414.
- [Che79] Jeff Cheeger, *Analytic torsion and the heat equation*, Annals of Mathematics **109** (1979), no. 2, 259–321.
- [CM10] Sylvain E Cappell and Edward Y Miller, *Complex-valued analytic torsion for flat bundles and for holomorphic bundles with $(1, 1)$ connections*, Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences **63** (2010), no. 2, 133–202.
- [de 36] Georges de Rham, *Sur les nouveaux invariants topologiques de M. Reidemeister.*, Rec. Math. Moscou, n. Ser. **1** (1936), 737–742.
- [Dei95] A. Deitmar, *Higher torsion zeta functions*, Advances in Mathematics **110** (1995), no. 1, 109 – 128.
- [DGRS19] Nguyen Viet Dang, Colin Guillarmou, Gabriel Riviere, and Shu Shen, *The Fried conjecture in small dimensions*, Inventiones mathematicae (2019), 1–55.
- [Fra35] Wolfgang Franz, *Über die torsion einer Überdeckung.*, Journal für die reine und angewandte Mathematik **173** (1935), 245–254.
- [Fri86] David Fried, *Analytic torsion and closed geodesics on hyperbolic manifolds*, Invent. Math. **84** (1986), no. 3, 523–540.
- [Fri87] ———, *Lefschetz formulas for flows*, Contemp. Math **58** (1987), 19–69.
- [Friedrich00] Thomas Friedrich, *Dirac operators in Riemannian geometry*, Graduate Studies in Mathematics, vol. 25, American Mathematical Society, Providence, RI, 2000, Translated from the 1997 German original by Andreas Nestke.
- [Gil84] Peter B. Gilkey, *The eta invariant and secondary characteristic classes of locally flat bundles*, Algebraic and differential topology—global differential geometry, Teubner-Texte Math., vol. 70, Teubner, Leipzig, 1984, pp. 49–87. MR 792686
- [GKM68] D. Gromoll, W. Klingenberg, and W. Meyer, *Riemannsche Geometrie im Grossen*, Lecture Notes in Mathematics, No. 55, Springer-Verlag, Berlin, 1968.

- [GS95] Gerd Grubb and Robert T. Seeley, *Weakly parametric pseudodifferential operators and Atiyah-Patodi-Singer boundary problems*, Invent. Math. **121** (1995), no. 3, 481–529.
- [LM89] H. Blaine Lawson, Jr. and Marie-Louise Michelsohn, *Spin geometry*, Princeton Mathematical Series, vol. 38, Princeton University Press, Princeton, NJ, 1989.
- [Mar88] A. S. Markus, *Introduction to the spectral theory of polynomial operator pencils*, Translations of Mathematical Monographs, vol. 71, American Mathematical Society, Providence, RI, 1988, Translated from the Russian by H. H. McFaden, translation edited by Ben Silver, with an appendix by M. V. Keldysh.
- [Mia80] Roberto J. Miatello, *The Minakshisundaram-Pleijel coefficients for the vector-valued heat kernel on compact locally symmetric spaces of negative curvature*, Trans. Amer. Math. Soc. **260** (1980), no. 1, 1–33.
- [Mil66] John Milnor, *Whitehead torsion*, Bulletin of the American Mathematical Society **72** (1966), no. 3, 358–426.
- [Mil78] John J. Millson, *Closed geodesics and the η -invariant*, Annals of Mathematics **108** (1978), no. 1, 1–39.
- [MM63] Yozô Matsushima and Shingo Murakami, *On vector bundle valued harmonic forms and automorphic forms on symmetric riemannian manifolds*, Ann. of Math. (2) **78** (1963), 365–416.
- [MM13] Simon Marshall and Werner Müller, *On the torsion in the cohomology of arithmetic hyperbolic 3-manifolds*, Duke Math. J. **162** (2013), no. 5, 863–888.
- [MS91] Henri Moscovici and Robert J. Stanton, *R-torsion and zeta functions for locally symmetric manifolds*, Inventiones mathematicae **105** (1991), no. 1, 185–216.
- [MSF18] Henri Moscovici, Robert J. Stanton, and Jan Frahm, *Holomorphic torsion with coefficients and geometric zeta functions for certain hermitian locally symmetric manifolds*, arXiv preprint arXiv:1802.08886 (2018).
- [Mül78] Werner Müller, *Analytic torsion and R-torsion of Riemannian manifolds*, Advances in Mathematics **28** (1978), no. 3, 233–305.
- [Mül11] ———, *A Selberg trace formula for non-unitary twists*, Int. Math. Res. Not. Imrn (2011), no. 9, 2068–2109.
- [Mül12] ———, *The asymptotics of the Ray-Singer analytic torsion of hyperbolic 3-manifolds.*, Metric and differential geometry. The Jeff Cheeger anniversary volume. Selected papers based on the presentations at the international conference on metric and differential geometry, Tianjin and Beijing, China, May 11–15, 2009, Berlin: Springer, 2012, pp. 317–352.

- [Par09] Jinsung Park, *Analytic torsion and Ruelle zeta functions for hyperbolic manifolds with cusps*, Journal of Functional Analysis **257** (2009), no. 6, 1713–1758.
- [Pfa12] Jonathan Pfaff, *Selberg and Ruelle zeta functions and the relative analytic torsion on complete odd-dimensional hyperbolic manifolds of finite volume*, Ph.d thesis, Bonn, 2012.
- [Pfa14] ———, *Analytic torsion versus Reidemeister torsion on hyperbolic 3-manifolds with cusps*, Mathematische Zeitschrift **277** (2014), no. 3-4, 953–974.
- [Pfa15] ———, *Selberg zeta functions on odd-dimensional hyperbolic manifolds of finite volume*, Journal für die Reine und Angewandte Mathematik (Crelles Journal) **2015** (2015), no. 703, 115–145.
- [Rei35] Kurt Reidemeister, *Homotopieringe und linsenräume*, Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg, vol. 11, Springer, 1935, pp. 102–109.
- [RR98] Yu B Rudyak and Julij Rudjak, *On thom spectra, orientability, and cobordism*, Springer Science & Business Media, 1998.
- [RS71] D. B. Ray and I. M. Singer, *R-torsion and the Laplacian on Riemannian manifolds*, Advances in Math. **7** (1971), 145–210.
- [See67] Robert T Seeley, *Complex powers of an elliptic operator*, Proc. Symp. Pure Math., vol. 10, 1967, pp. 288–307.
- [She18] Shu Shen, *Analytic torsion, dynamical zeta functions, and the Fried conjecture*, Analysis and PDE **11** (2018), no. 1.
- [Spi15] Polyxeni Spilioti, *Functional equations of Selberg and Ruelle zeta functions for non-unitary twists*, arXiv preprint arXiv:1507.05947 (2015).
- [Spi18] ———, *Selberg and Ruelle zeta functions for non-unitary twists*, Annals of Global Analysis and Geometry **53** (2018), no. 2, 151–203.
- [Wal60] CTC Wall, *Determination of the cobordism ring*, Annals of Mathematics (1960), 292–311.
- [Wal76] Nolan R. Wallach, *On the Selberg trace formula in the case of compact quotient*, Bull. Amer. Math. Soc. **82** (1976), no. 2, 171–195.
- [Wot08] A. Wotzke, *Die Ruellesche Zetafunktion und die analytische Torsion hyperbolischer Mannigfaltigkeiten*, Ph.d thesis, Bonn, Bonner Mathematische Schriften, 2008.

P. SPILIOTI, Department of Mathematics, Ny Munkegade 118, 8000, Aarhus C, Denmark
E-mail address: spilioti@math.au.dk